



Using what we know about threat reactivity models to understand mental health during the COVID-19 pandemic

Carter J. Funkhouser^{a,b,*}, David M. Klemballa^a, Stewart A. Shankman^a

^a Northwestern University, Department of Psychiatry and Behavioral Sciences, 680 N. Lake Shore Dr., Chicago, IL, 60611, USA

^b University of Illinois at Chicago, Department of Psychology, 1007 W. Harrison St., Chicago, IL, 60607, USA

ARTICLE INFO

Keywords:

Stress
Uncertainty
Psychopathology
Anxiety
Fear
COVID-19

ABSTRACT

The COVID-19 pandemic has been accompanied by unprecedented levels of stress and threats in a variety of domains (e.g., health, livelihood). Individual differences in threat reactivity may explain why some individuals are at elevated risk for the development or maintenance of psychopathology during the COVID-19 pandemic. This article describes several prominent models, mechanisms, and components of threat reactivity (e.g., appraisals, intolerance of uncertainty, avoidance) and discusses how they might help improve understanding of changes in psychopathology during and following the COVID-19 pandemic.

The COVID-19 pandemic is an unprecedented global public health crisis. Hundreds of millions of people worldwide have been infected with COVID-19 and millions have died and continue to die from this unrelenting virus. Furthermore, many COVID-19 survivors experience a post-COVID-19 condition (often termed “long COVID”; Huang et al., 2021) involving prolonged symptoms (e.g., fatigue, shortness of breath, cognitive dysfunction) that last for at least 2 months and cannot be explained by an alternative diagnosis (Soriano, Murthy, Marshall, Relan, & Diaz, 2021). The pandemic has also been associated with increased rates of suicidal thoughts and behaviors (Dubé, Smith, Sherry, Hewitt, & Stewart, 2021) and intimate partner violence (Richards, Nix, Mourtgos, & Adams, 2021). Researching the mental health consequences of the COVID-19 pandemic is an urgent research priority (Holmes et al., 2020). Preliminary studies suggest that psychological distress and mental health symptoms generally increased in the early months of the pandemic (e.g., March–April 2020), largely returned to pre-pandemic levels in the ensuing months (Aknin et al., 2021; Robinson, Sutin, Daly, & Jones, 2021; Santomauro et al., 2021), and have wavered during the arrival of new variants of the virus (Cohen, 2021). However, there has been substantial heterogeneity in the mental health consequences of COVID-19 (Gloster et al., 2020) and it is important to understand individual differences that may increase risk for psychopathology during the COVID-19 pandemic.

Individual differences in *threat reactivity* may play a critical role in identifying who might be at particular risk for mental health difficulties during and following the COVID-19 pandemic. Threat reactivity refers

to a cross-species set of processes and sensitivities that have been implicated in numerous internalizing and externalizing psychopathologies, most notably post-traumatic stress disorder, substance use, and various anxiety disorders (Lieberman et al., 2020; Stevens et al., 2021; Williams et al., 2016). The transdiagnostic nature and clinical importance of threat reactivity has made it a particularly important set of factors/mechanisms given the increased interest in alternative psychiatric nomenclatures proposed in recent years –e.g., NIMH’s Research Domain Criteria (RDoC; Funkhouser et al., 2021; Zalta & Shankman, 2016), the Hierarchical Taxonomy of Psychopathology (Kotov et al., 2017), and the ability to study (and ultimately treat) threat reactivity across multiple units of analyses (Shankman & Gorka, 2015; Young et al., 2021). The goal of this paper is therefore to describe the role that different aspects and mechanisms of threat reactivity may play in the occurrence or risk for psychopathology during and/or following the COVID-19 pandemic.

Several “disclaimers” and caveats should be noted. First, the purpose of this paper is *not* to argue that threat reactivity is universally maladaptive. Like most psychological constructs, individual differences in threat reactivity are likely to be normally distributed. A *certain amount* of heightened threat reactivity (and its emotional concomitants, fear and anxiety; Daniel-Watanabe & Fletcher, 2021; Klumpp & Shankman, 2018) during COVID-19 is a healthy response to a worldwide pandemic as it promotes adaptive safety behaviors such as hand washing, social distancing, and mask wearing (Harper, Satchell, Fido, & Latzman, 2020; Knowles & Olatunji, 2021). However, both excessive *and* insufficient

* Corresponding author. University of Illinois at Chicago, Department of Psychology, 1007 W. Harrison St., Chicago, IL, 60607, USA.

E-mail address: carterfunkhouser@gmail.com (C.J. Funkhouser).

levels of threat reactivity may lead to maladaptive behaviors in response to COVID. For example, low levels of threat reactivity may, in part, lead to behaviors that increase risk for COVID-19 infection and transmission, and high levels of threat reactivity might lead to excessive distress and maladaptive risk-averse behaviors (e.g., refusal to leave home; excessive handwashing). In other words, the ‘right amount’ of threat reactivity is adaptive – analogous to the evolutionary advantages of reacting with fear if confronted with a present and unambiguous threat (Lang, 2010).

Second, COVID-19 is associated with different types of threats, including catching the virus oneself, having a loved one catch the virus, economic consequences, and social isolation (Coelho, Suttiwan, Arato, & Zsido, 2020). These threats also likely played different roles at different stages of the pandemic given the ever-changing landscape of case rates, mitigation strategies, virus variants, etc. Relatedly, there have been other sources of stress concurrent with the pandemic (e.g., political and racial tensions, protests) that have contributed to the mental health crisis during the last two years. Distinguishing the sources of threats/stress during the COVID-19 pandemic is beyond the scope of this paper and will be addressed, in part, by other articles in this special issue (see McLaughlin article). This paper will therefore only focus on within-individual mechanisms, processes, and features of threat reactivity and how they may play a role in COVID-related psychopathology. However, it is important to note that these within-person aspects of threat reactivity likely interact with particular sources of threat/stress in different ways in predicting psychopathology, thus creating different *vulnerability-stress interactions* (Monroe & Simons, 1991).

Third, threat reactivity may change as a result of chronic stress exposure during the COVID-19 pandemic (see McLaughlin et al., this issue for further discussion of mechanisms linking chronic stress to psychopathology). The effects of chronic or repeated stressors on threat reactivity specifically are poorly understood. Many of the aspects of threat reactivity discussed in this paper have been largely studied by averaging threat responses elicited during relatively short laboratory paradigms (Denson, Spanovic, & Miller, 2009; Duits et al., 2015; Grillon, 2002). Examining the *time course* of threat reactivity (a series of constructs often referred to as affective chronometry; Davidson, 1998) during these laboratory paradigms may provide insights into the impacts of chronic stressors on threat reactivity (Klumpp & Shankman, 2018). However, the extent to which these laboratory paradigms generalize to the chronic stressors commonly experienced during the COVID-19 pandemic is unclear.

1. Different aspects of threat reactivity could lead to adverse mental health outcomes during COVID-19

1.1. Threat appraisals

Appraisal, or the process of detecting and assessing environmental features with regard to one’s well-being (Moors, Ellsworth, Scherer, & Frijda, 2013), is an important driver of emotions and behavior (Sheeran, Harris, & Epton, 2014). Threat appraisals evaluate (a) whether a stimulus or situation is potentially harmful to one’s well-being, (b) the likelihood of the dangerous or negative event, (c) the severity of the event if it were to occur, and (d) the ability to cope and access resources if necessary (Carr, 1974; Lazarus, 1991; Salkovskis, 1996). Individuals with anxiety symptoms or disorders tend to display biases in each of these domains such that they tend to overestimate the likelihood of a negative event, overestimate the negative impact of the event if it were to happen, and underestimate the availability of coping resources (Salkovskis, 1996). Cognitive behavioral interventions for anxiety disorders are thought to reduce anxiety by helping patients develop more realistic and adaptive threat appraisals (Clark & Beck, 2010), and empirical studies suggest that cognitive behavioral therapy improves threat reappraisal (Draheim & Anderson, 2021), which may mediate reduction in anxiety symptoms (Smits, Julian, Rosenfield, & Powers, 2012).

Threat appraisals have played an important role in shaping thoughts,

emotions, and behaviors related to COVID-19. Greater appraisals of the risk and/or severity of contracting COVID-19 have been associated with greater COVID-19-related fear and distress (Mertens, Gerritsen, Duijndam, Saleminck, & Engelhard, 2020; Taylor, Landry, Paluszek, & Asmundson, 2020), intention to get vaccinated (Caserotti et al., 2021), and adherence to social distancing and avoidance of public places (Al-Hasan, Khuntia, & Yim, 2020; Taylor et al., 2020). These findings corroborate data from previous disease outbreaks suggesting that greater perceptions of the likelihood and/or severity of infection were associated with pandemic-related anxiety and safety behaviors (Rudisill, 2013). Importantly, COVID-19-related worry and protective behaviors are more strongly predicted by *perceived* COVID-19 threat than *actual* COVID-19 threat (Schmidt et al., 2021), highlighting the importance of threat appraisals in shaping emotions and behavior during the pandemic. In addition to perceived health-related threats, perceived risk of suffering personal economic losses due to COVID-19 is associated with preventative health behaviors and support for policies to mitigate COVID-19 across the globe (Nisa et al., 2021).

Considering the influence of threat perceptions on COVID-19 protective behaviors (e.g., social distancing, wearing masks, vaccination), it is worth examining how COVID-19 threat perceptions are formed. Of course, the presence or absence of known risk factors for contracting COVID-19 (e.g., number of cases in one’s community) or experiencing severe COVID-19 symptoms (e.g., having chronic medical conditions) influence many people’s COVID-19 threat perceptions. These known risk factors are only moderately correlated with COVID-19 risk perceptions, however (e.g., $b = 0.24$; Ahuja et al., 2021). The COVID-19 pandemic has been accompanied by an overabundance of information and misinformation (Naeem & Bhatti, 2020), and risk perceptions are generally influenced by what information is most salient or available (Tversky & Kahneman, 1973). Information and misinformation have been most commonly received from government agencies or officials, news media, social media, family, and/or friends. Interestingly, several studies have shown that trust in government, science, and medical professionals as information sources was associated with greater COVID-19 risk perception and greater COVID-19 knowledge (e.g., Roozenbeek et al., 2020), whereas individuals whose most trusted COVID-19 information source was other news sources or social media were less knowledgeable about COVID-19 (Sakya et al., 2021). Relatedly, the politicization of the COVID-19 pandemic and associated policies in the United States has likely influenced the amount of trust placed in government officials and agencies relative to other information sources. Indeed, political affiliation was associated with COVID-19 risk perceptions and protective behaviors in the United States (Bruine de Bruin, Saw, & Goldman, 2020), and individuals may be more likely to trust government officials whose political affiliation matches their own. Reliance on unreliable information sources is likely to continue due to confirmation bias (i.e., the tendency to favor information in a way that supports one’s prior beliefs; Nickerson, 1998), thereby minimizing opportunities to generate more accurate reappraisals of COVID-19 threats.

1.2. Attentional bias

Attentional bias to threat, or the preferential allocation of attention to threat-related stimuli over neutral stimuli, is adaptive for the early detection of threats. However, attentional bias to threat is heightened in individuals with or at risk for anxiety disorders (Bar-Haim, Lamy, Pergamin, Bakermans-Kranenburg, & Van Ijzendoorn, 2007), and numerous cognitive theories implicate elevated attentional bias to threat in the etiology of fear- and anxiety-related psychopathology (e.g., Beck & Clark, 1997). For instance, individuals high in trait anxiety may exhibit vigilance toward threat during an early stage of immediate threat processing, followed by a later attentional allocation stage characterized by avoidance (Cisler & Koster, 2010).

There is preliminary evidence that, on average, individuals have exhibited attentional bias to COVID-19-related stimuli relative to

neutral stimuli during the COVID-19 pandemic (Albery, Spada, & Nikčević, 2021; Cannito et al., 2020). More importantly, COVID-19-related attentional bias was associated with more frequent symptoms of COVID-related anxiety (Albery et al., 2021) and health anxiety (Cannito et al., 2020), suggesting that attentional bias to threat may be involved in the development of anxiety specifically regarding the COVID-19 pandemic.

Assessing, and ultimately intervening on, attentional bias is likely to require novel methods given that traditional methods of assessing attentional bias index have questionable psychometric properties and often conflate multiple distinct processes, e.g., orientation toward threat (i.e., vigilance) and disengagement from threat (Koster, Crombez, Verschuere, & De Houwer, 2004; MacLeod, Mathews, & Tata, 1986; Pettit et al., 2020). Metrics aiming to capture distinct attentional bias components may have superior psychometric properties to traditional metrics (Evans, Walukevich, Seager, & Britton, 2018), are elevated in individuals with or at risk for anxiety disorders (Evans, Walukevich, & Britton, 2016; Meissel et al., 2021), and may be differentially associated with COVID-related anxiety (Albery et al., 2021; Nasiry & Ameli, 2021). These findings from laboratory studies of attentional bias are generally consistent with results from self-report studies. Greater COVID-19 news exposure was associated with increased state anxiety (Nekliudov et al., 2020) and depressive symptoms (Olagoke, Olagoke, & Hughes, 2020), and individuals reporting greater threat-related cognitive biases perceived COVID-19 infection as being more likely and were more likely to use maladaptive emotion regulation strategies (e.g., suppression; Schudy, Żurek, Wiśniewska, Piejka, Gawęda, & Okruszek, 2020).

1.3. Avoidance

Avoidant behavior is characterized as a conscious or unconscious effort to avoid a stressor or threat, often to minimize negative thoughts and emotions brought on by the stressor (Barlow, 2004). As a behavioral response, although effective in the short-term (but see Kryptos, Eftting, Kindt, & Beckers, 2015), avoidance serves to maintain fears and worries, often resulting in decreased well-being and onset of depression and anxiety disorders (Hofmann & Hay, 2018; Kryptos et al., 2015). The COVID-19 pandemic reflects significant stressors that interact with avoidant responses, resulting in decreased psychological well-being.

Avoidance behaviors can lead to psychopathology during COVID-19 through multiple pathways. Many people have gone to significant lengths to avoid exposure to the virus by significantly limiting their pre-pandemic activities (e.g., exercising, socializing with friends). While avoiding these activities can be adaptive in reducing COVID-19 risk, they can also increase risk for psychopathology by removing adaptive coping strategies and sources of social support (Solomou & Constantinidou, 2020). Another pathway through which avoidant behaviors can lead to psychopathology is when individuals disregard the threat posed by the virus (i.e., “disengagement coping”; Skinner, Edge, Altman, & Sherwood, 2003; Taha, Matheson, Cronin, & Anisman, 2014), as this could lead to maladaptive avoidance of unpleasant feelings related with the pandemic. “Disengagement coping” could also account for the significant increased frequency of binge drinking during COVID-19, findings that are especially pronounced in those with depression (Greenglass, Chiacchia, & Fiskensbaum, 2021; Weerakoon, Jetelina, & Knell, 2021). Avoiding recognition of the seriousness of the threat of COVID-19 is also risky as this can lead to unsafe behavior that can contribute to the virus spread (e.g., socializing in close physical proximity to others, refusal to wear masks in public spaces, neglecting handwashing). In sum, avoidance behaviors are multifaceted and identifying the function of the specific individual’s behaviors is of the utmost importance (Hofmann & Hay, 2018). Moreover, given the transmissibility of COVID-19, avoidance behaviors likely require a “goldilocks” approach to identify the “just right” amount of avoidance behaviors.

1.4. Fear acquisition and extinction

Rachman’s (1977) three-pathway model of fear acquisition suggests that fear can be acquired through either fear learning (i.e., the process by which a neutral conditioned stimulus elicits fear over time following repeated pairing with an aversive unconditioned stimulus), information transmission (i.e., being informed that a stimulus is dangerous), or modeling (i.e., observing a fearful response to a stimulus in others). Fear learning has received the most attention as a potential mechanism of pathological fear (e.g., anxiety disorders) and can be elicited by classic Pavlovian conditioning paradigms. Individuals with an anxiety disorder display elevated conditioned responding during fear learning compared to healthy controls (Lissek et al., 2005), supporting its putative etiological role in phobic anxiety disorders (Field, 2006; Grillon, 2008). Patients with anxiety disorders also exhibit difficulty inhibiting a fear response when presented with a safety cue (Lissek et al., 2005), suggesting that fear overgeneralization or impaired safety learning may also be involved (Jovanovic, Kazama, Bachevalier, & Davis, 2012). As discussed previously, fear responses are often accompanied by behaviors (e.g., safety behaviors, avoidance) that temporarily reduce distress, but may contribute to the maintenance of fear in the long term (Lovibond, Davis, & O’Flaherty, 2000). On the other hand, fear extinction is the process by which a fear response gradually decreases following repeated exposure to a feared stimulus in the absence of an associated aversive event (Myers & Davis, 2007). Fear extinction is thought to be driven by inhibition learning (Craske et al., 2008) or habituation processes (Foa & Kozak, 1986), and is impaired in individuals with anxiety disorders compared to healthy controls (Duits et al., 2015; Lissek et al., 2005). As such, fear extinction is widely studied in the context of exposure therapy, which itself is a clinical proxy of fear extinction (Craske et al., 2008).

Widespread fear learning has taken place during the COVID-19 pandemic. Previously neutral stimuli (e.g., the word “coronavirus”) became conditioned stimuli and elicited a fear response (Presti, Mchugh, Gloster, Karekla, & Haye, 2020). There are a variety of aversive outcomes that might act as unconditioned stimuli in the context of COVID-19 (e.g., loss of employment, hospitalization, death of loved ones). Additionally, overgeneralized fear learning in the context of COVID-19 (e.g., exhibiting a fear response to a stimulus that is not associated with COVID-19 risk) may increase vulnerability for maladaptive COVID-19-related anxiety. As COVID-19 cases declined, many communities gradually reduced mitigation policies and allowed individuals to resume pre-pandemic activities and reduce or discontinue protective behaviors (e.g., mask-wearing, social distancing). This process resembles that of fear extinction. For example, if someone had previously avoided certain activities to minimize COVID-19 risk (e.g., dining in a restaurant) and gradually resumes those activities without experiencing any negative outcomes, they may form new beliefs about the likelihood of negative COVID-related outcomes associated with those activities. Impaired inhibitory learning could lead to the maintenance of COVID-19 fear even as COVID-19 becomes less threatening (i.e., poor inhibitory learning still prevents them from dining out in restaurants). Of course, negative outcomes may still occur when COVID-19-related threats are decreasing, which could induce rapid fear reacquisition regarding COVID-19-related threats (Woods & Bouton, 2007). Indeed, many parts of the world have experienced substantial fluctuations in COVID-19 cases and associated threats and preventative policies. Calibrating one’s fear to changes in environmental threats is adaptive, and elevated fear responses during fear learning or impaired inhibitory learning during extinction may increase risk for excessive COVID-19 fear and associated mental health problems (Craske, Hermans, & Vervliet, 2018).

Although less studied than fear learning in the traditional anxiety literature, information transmission and modeling may also partially explain COVID-19 fear acquisition. For example, many people sought out information at the beginning of the pandemic to help them more

accurately assess COVID-19 threats and modify their behavior accordingly (Bento et al., 2020). Modeling has also likely played a role in shaping fears associated with COVID-19, especially considering that behaviors are more likely to be influenced by modeling in times of uncertainty (Smith, Hogg, Martin, & Terry, 2007). For instance, the phenomenon of ‘panic buying’ (i.e., excessively stockpiling food and goods) in the initial stage of the pandemic may have resulted from many people modeling a relatively small group of over-anxious, high IU individuals (Arafat et al., 2020; Taylor, 2021). Additionally, consistent with modeling playing a role in COVID-19 fear acquisition, college students’ perceptions of their peers’ adherence to preventative guidelines and COVID-19 vaccination intentions were positively associated with their own behaviors and vaccination intentions (Graupensperger, Abdallah, & Lee, 2021; Graupensperger, Lee, & Larimer, 2021). The impact of modeling is also evidenced by the role of celebrities and other public figures in encouraging (or discouraging) COVID mitigation strategies (Martinez-Berman, McCutcheon, & Huynh, 2021).

1.5. Neural correlates of threat reactivity

There is a large literature of human and animal studies examining the neuroanatomy and neural circuitry involved in threat reactivity in clinical samples and in healthy populations. An extensive discussion of this literature is beyond the scope of this article and has been reviewed in other papers (e.g., Grupe & Nitschke, 2013), some of which focus on a specific component of threat processing such as fear learning (Ohman & Mineka, 2001), extinction learning (Myers & Davis, 2007), or threat attention and appraisal (Britton, Lissek, Grillon, Norcross, & Pine, 2011). Broadly, threat responses are generated in regions in the limbic system (e.g., the amygdala) and modulated by areas such as the prefrontal cortex (Marek, Strobel, Bredy, & Sah, 2013). Other regions have been implicated in reactivity to certain kinds of threats. For example, the anterior insula is involved in the anticipatory processing of uncertain threats (Shankman et al., 2014; Walker, Toufexis, & Davis, 2003). Functional neuroimaging studies suggest that individuals with anxiety and traumatic disorders often show amygdala and anterior insula hyperactivation and hypoactivation in prefrontal regions involved in emotion modulation (Etkin & Wager, 2007).

Few (if any) neuroimaging studies to date have examined neural reactivity to threatening stimuli specifically related to COVID-19. However, several studies have examined whether neural reactivity to other threats (e.g., fearful faces) assessed before the COVID-19 pandemic predicted mental health outcomes during the pandemic. For example, pre-pandemic amygdala activation to threat predicted internalizing symptoms during the pandemic and moderated the association between COVID-19-related stressors and internalizing symptoms (Weissman et al., 2021). Additionally, higher prefrontal activity during emotion regulation predicted lower stress burden during the COVID-19 pandemic (Monninger et al., 2021) and anterior insula reactivity to uncertain threat predicted negative affect during the COVID-19 pandemic (Khorrami, Manzler, Kreutzer, & Gorka, 2021). Taken together, these studies suggest that neural markers of threat reactivity may play a prognostic role in COVID-19-related psychopathology.

1.6. Neuroendocrine correlates of threat reactivity

The neuroendocrine system plays a central role in stress and emotion (dys)regulation (McEwen & Akil, 2020) and dysregulation in this system has been implicated in a variety of psychological disturbances including depression, PTSD, and anxiety (Ehlert, Gaab, & Heinrichs, 2001). Within the neuroendocrine system, acute or chronic stressors induce a psychological and physiological response to maintain homeostasis, which activates the hypothalamic-pituitary-adrenal (HPA) axis. A detailed description of the functioning of the HPA axis is beyond the scope of this paper, though it is important to note that this system is responsible for secreting cortisol which ultimately operates as a negative feedback

mechanism within this system (Stokes, 1995). Subjective threat appraisals influence the neuroendocrine response (Schlotz, Hammerfal, Ehlert, & Gaab, 2011). Accordingly, threat appraisals may impact the neuroendocrine system and its role in emotion regulation during COVID-19.

COVID-19 threats have also led many to self-quarantine during the pandemic, thus increasing social isolation and feelings of loneliness (Hoffart, Johnson, & Ebrahimi, 2020). Loneliness is associated with neuroendocrine abnormalities (e.g., HPA axis activation; Doane & Adam, 2010) and internalizing psychopathologies (Shevlin, McElroy, & Murphy, 2015). Together, this suggests that the self-quarantine behavior associated with virus-related threat detection may contribute to a neuroendocrine response that underlies emotional dysregulation and potential psychopathology.

1.7. Intolerance of uncertainty

Another critical component of threat reactivity is how individuals respond to uncertainty. Uncertainty itself can be perceived as threatening (Dugas, Letarte, Rhéaume, Freeston, & Ladouceur, 1995), even in the context of reward (Nelson, Shankman, & Proudfit, 2014). Individuals with elevated intolerance of uncertainty (IU) – the dispositional inability to endure the aversive response triggered by perceived uncertainty (Carleton, 2016) – are more likely to experience anxiety and worry and engage in uncertainty-reducing behaviors in response to uncertainty (Badia, Harsh, & Abbott, 1979; Freeston, Tiplady, Mawn, Bottesi, & Thwaites, 2020; Ladouceur, Gosselin, & Dugas, 2000). IU is associated with increased risk for anxiety and depressive disorders (Correa, Liu, & Shankman, 2019; Funkhouser et al., 2021; Gentes & Ruscio, 2011) and may connote vulnerability for psychopathology due to increased contextual uncertainty during COVID-19. IU also may be a process of change in multiple cognitive behavioral psychotherapies. IU decreases during cognitive behavioral therapy (CBT) and reductions in IU are associated with reductions in anxiety disorder symptoms (Khakpoor, Mohammadi Bytamar, & Saed, 2019; Mahoney & McEvoy, 2012; McEvoy & Erceg-Hurn, 2016), suggesting that IU may be a good target for psychotherapy during the COVID-19 pandemic (Dugas & Ladouceur, 2000; van der Heiden, Muris, & van der Molen, 2012).

Preliminary studies conducted during the COVID-19 pandemic suggest that IU was associated with greater health anxiety, pandemic-related anxiety, appraisals of pandemic-related threat severity, use of maladaptive emotion-focused coping strategies, and conspiratorial thinking (Larsen, Donaldson, Liew, & Mohanty, 2021; Mertens et al., 2020; Satici, Saricali, Satici, & Griffiths, 2020; Tull et al., 2020). These results corroborate findings from a study conducted during the H1N1 pandemic (Taha, Matheson, Cronin, & Anisman, 2014). IU has also been associated with uncertainty-reducing behaviors related to health anxiety such as seeking medical information online and ‘panic buying’ or stockpiling supplies (Taylor, 2021). Although information-seeking regarding COVID-19 can be helpful for informing personal choices, frequent consumption of COVID-related news was associated with greater uncertainty (Yoon et al., 2021) and can exacerbate worry and anxiety (Soroya, Farooq, Mahmood, Isoaho, & Zara, 2021). The effect of information-seeking on COVID-19 fear and anxiety may be especially strong in individuals with high IU (Baerg & Bruchmann, 2022; Bottesi, Marino, Vieno, Ghisi, & Spada, 2021). For example, suppose a person high in IU is worried that they had contracted COVID-19 because they experienced a headache and sore throat. To attempt to reduce uncertainty, they may search for information about the likelihood of this outcome on the internet, where they would likely encounter a variety of diagnostic and prognostic explanations and probabilities, which will further trigger their intolerance of uncertainty. This would result in an escalation in worry and health anxiety (Fergus, 2013), a phenomenon termed ‘cyberchondria’ (White & Horvitz, 2009). Consistent with this vicious cycle, recent studies found that IU partially accounted for the associations between health anxiety/obsessive compulsive symptoms

and fear of COVID-19 (Wheaton, Messner, & Marks, 2021) and between anxiety-related coping styles and compulsive buying behavior (Çelik & Köse, 2021).

Interestingly, the temporal course of COVID-19-related uncertainty coincided with the changes in symptoms of internalizing disorders (e.g., Major Depression, anxiety disorders) during the COVID-19 pandemic. The early months of the COVID-19 pandemic in the spring of 2020 were characterized by salient yet poorly understood (i.e., uncertain) threats, and this increase in uncertainty may have been involved in the development of internalizing symptoms in individuals high in IU during this time. Some COVID-related threats (e.g., COVID-19 infection) have become better understood over the course of the pandemic, thereby reducing uncertainty. On the other hand, there is ongoing uncertainty regarding other threats associated with COVID-19. For example, uncertainty regarding COVID-19 vaccines' effectiveness and potential adverse effects have been cited as reasons for vaccine hesitancy (Solís Arce et al., 2021). Individuals high in IU may be especially distressed by this ongoing elevated uncertainty, leading to especially high levels of psychopathology in these individuals. That said, IU may not impact everyone's beliefs about COVID-19. For instance, some individuals may be 100% certain in their beliefs related to COVID-19 (e.g., regarding vaccine efficacy), and IU would have little impact on mental health for these individuals.

2. How different aspects of threat reactivity might work together

Aspects of threat reactivity have largely been discussed separately thus far, but threat-related psychopathology likely arises from complex and dynamic relationships involving multiple aspects of threat reactivity (Fried & Robinaugh, 2020; Robinaugh et al., 2019). Understanding how processes such as these relate to each other both within and across biobehavioral systems (e.g., neural circuits, behavior) is a central goal of the RDoC initiative (Insel et al., 2010) and has been examined in numerous studies. For example, if an individual engages in adaptive threat reappraisals, this can accelerate extinction learning and reduce threat-related attentional bias (Blechert et al., 2015; Van Damme, Crombez, Hermans, Koster, & Eccleston, 2006). Alternatively, avoidance of feared situations prevents opportunities for extinction learning and threat reappraisal (Craske et al., 2018; Lovibond, Mitchell, Minard, Brady, & Menzies, 2009). In fact, studies have shown that individuals may infer danger from the use of avoidance behaviors even when the avoided situations are nonthreatening (Blakey & Abramowitz, 2016; Engelhard, van Uijen, van Seters, & Velu, 2015; van Uijen, Leer, & Engelhard, 2018). This in turn may reinforce potentially inaccurate threat appraisals and contribute to the persistence of attentional biases and fear (Britton et al., 2011; Heeren & McNally, 2016).

Individuals with high IU may be especially reactive to threats. In uncertain situations, individuals with high IU are prone to greater threat appraisals (Pepperdine, Lomax, & Freeston, 2018), avoidance (Flores, López, Vervliet, & Cobos, 2020; San Martín, Jacobs, & Vervliet, 2020), and neural reactivity to threat in regions including the amygdala and anterior insula (Tanovic, Gee, & Joermann, 2018). High IU is also associated with poorer threat extinction learning (Morriss, Wake, Elizabeth, & van Reekum, 2021; Morriss, Zuj, & Mertens, 2021), greater sensitivity to instructions about safety (Gorka, Lieberman, Nelson, Sarapas, & Shankman, 2014; Mertens & Morriss, 2021; Morriss, Bennett, & Larson, 2021; Morriss & van Reekum, 2019), greater threat generalization (Bauer et al., 2020), and heightened attentional biases to uncertainty (Morriss, McSorley, & van Reekum, 2018; Morriss & McSorley, 2019). These findings collectively suggest that individuals with high IU may exhibit deficits in other aspects of threat reactivity and thus be more sensitive to COVID-19 threats and struggle to adjust to changing information about COVID-19 threats and safety guidelines.

3. Conclusions

The COVID-19 pandemic has increased contextual threats worldwide and mechanisms and aspects of threat reactivity may help explain why some individuals were (and are) at increased risk for developing mental health problems during this time. Some of these threat processes may be more distal risk factors for psychopathology, whereas others are thought to be more proximally involved in etiology. Additionally, although discussed independently above, the different components and mechanisms of threat reactivity likely interact and relate in important ways (e.g., appraisals affecting avoidance behaviors; Hofmann & Hay, 2018). A common theme across threat reactivity models is that excessive threat reactivity may contribute to the development of mental health problems such as maladaptive anxiety. On the other hand, insufficient threat reactivity may lead to behaviors that increase risk for negative outcomes related to COVID-19.

Fortunately, processes involved in threat reactivity can be reduced through evidence-based interventions (e.g., exposure therapy, selective serotonin reuptake inhibitors) and federal governments in many countries have allocated funding for expanding access to mental health resources (e.g., the Consolidated Appropriations Act, 2021 in the United States). Increasing the scalability of these interventions will be important to support wider dissemination.

It is also worth noting that existing mental health interventions largely aim to reduce threat reactivity, but some individuals underestimate COVID-19 threats and might benefit from increased COVID-19 threat reactivity. Although comparatively less likely to contribute to internalizing mental health problems, underestimations of COVID-19 threats are associated with less engagement in recommended protective behaviors (Taylor et al., 2020) and may interfere with ongoing efforts to mitigate COVID-19 transmission. Increasing threat perceptions regarding COVID-19 remains an ongoing public health challenge (Ali, 2020). Many interventions occur at the level of the individual, but societal-level interventions (e.g., clear and consistent messaging) can also effectively modulate COVID-19 threat reactivity. Individuals with maladaptively low COVID-19 threat perceptions tend to distrust government officials and scientists (Roozenbeek et al., 2020), and delivering interventions through alternative, more trusted information sources may be more effective for modifying this group's COVID-19 threat perceptions and related behavior.

In sum, the 'right amount' of threat reactivity is adaptive for detecting and responding to threats related to COVID-19, but insufficient or excessive levels of COVID-19 threat reactivity can increase risk for mental or physical health problems. Interventions that help individuals 'calibrate' their COVID-19 threat sensitivity in accordance with available (and constantly changing) information about COVID-19 are critical for preventing these negative outcomes.

Funding

This work was supported by National Institute of Mental Health grants R42127971 (PI: Shankman) and F31MH123042 (PI: Funkhouser).

CRedit authorship contribution statement

Carter J. Funkhouser: Conceptualization, Writing – original draft, Writing – review & editing. **David M. Klemballa:** Writing – original draft, Writing – review & editing. **Stewart A. Shankman:** Conceptualization, Writing – original draft, Writing – review & editing.

Declaration of competing interest

None.

References

- Ahuja, M., Mamudu, H. M., Weierbach, F. M., Dowling-McClay, K., Stewart, D. W., Awasthi, M., et al. (2021). Perceptions of risk for COVID-19 among individuals with chronic diseases and stakeholders in Central Appalachia. *Humanities and Social Sciences Communications*, 8(1), 1–6. <https://doi.org/10.1057/s41599-021-00906-7>
- Aknin, L., Neve, J.-E. De, Dunn, E., Fancourt, D., Goldberg, E., Helliwell, J., et al. (2021). *Mental health during the first year of the COVID-19 pandemic: A review and recommendations for moving forward*. <https://doi.org/10.31234/OSF.IO/ZW93G>. Perspectives on Psychological Science.
- Al-Hasan, A., Khuntia, J., & Yim, D. (2020). Threat, coping, and social distance adherence during COVID-19: Cross-continental comparison using an online cross-sectional survey. *Journal of Medical Internet Research*, 22(11), Article e23019. <https://doi.org/10.2196/23019>
- Albery, I. P., Spada, M. M., & Nikčević, A. V. (2021). The COVID-19 anxiety syndrome and selective attentional bias towards COVID-19-related stimuli in UK residents during the 2020–2021 pandemic. *Clinical Psychology & Psychotherapy*. <https://doi.org/10.1002/cpp.2639>
- Ali, S. (2020). *Combatting against covid-19 & misinformation: A systematic review*. <https://doi.org/10.1007/s42087-020-00139-1>. *Human Arenas*.
- Arafat, S. M. Y., Kar, S. K., Marthoenis, M., Sharma, P., Hoque Apu, E., & Kabir, R. (2020). Psychological underpinning of panic buying during pandemic (COVID-19). *Psychiatry Research*, 289, 113061. <https://doi.org/10.1016/j.psychres.2020.113061>
- Badia, P., Harsh, J., & Abbott, B. (1979). Choosing between predictable and unpredictable shock conditions: Data and theory. *Psychological Bulletin*, 86(5), 1107–1131. <https://doi.org/10.1037/0033-2909.86.5.1107>
- Baerg, L., & Bruchmann, K. (2022). COVID-19 information overload: Intolerance of uncertainty moderates the relationship between frequency of internet searching and fear of COVID-19. *Acta Psychologica*, 224, 103534. <https://doi.org/10.1016/j.actpsy.2022.103534>
- Bar-Haim, Y., Lamy, D., Pergamin, L., Bakermans-Kranenburg, M. J., & Van Ijzendoorn, M. H. (2007). Threat-related attentional bias in anxious and nonanxious individuals: A meta-analytic study. *Psychological Bulletin*, 133(1), 1–24. <https://doi.org/10.1037/0033-2909.133.1.1>
- Barlow, D. H. (2004). *Anxiety and its disorders: The nature and treatment of anxiety and panic* (2nd ed.). Guilford Press.
- Bauer, E. A., MacNamara, A., Sandre, A., Lonsdorf, T. B., Weinberg, A., Morris, J., et al. (2020). Intolerance of uncertainty and threat generalization: A replication and extension. *Psychophysiology*, 57(5), e13546. <https://doi.org/10.1111/psyp.13546>
- Beck, A. T., & Clark, D. A. (1997). An information processing model of anxiety: Automatic and strategic processes. *Behaviour Research and Therapy*, 35(1), 49–58. [https://doi.org/10.1016/S0005-7967\(96\)00069-1](https://doi.org/10.1016/S0005-7967(96)00069-1)
- Bento, A. I., Nguyen, T., Wing, C., Lozano-Rojas, F., Ahn, Y.-Y., & Simon, K. (2020). Evidence from internet search data shows information-seeking responses to news of local COVID-19 cases. *Proceedings of the National Academy of Sciences*, 117(21), 11220–11222. <https://doi.org/10.1073/pnas.2005335117>
- Blakey, S. M., & Abramowitz, J. S. (2016). The effects of safety behaviors during exposure therapy for anxiety: Critical analysis from an inhibitory learning perspective. *Clinical Psychology Review*, 49, 1–15. <https://doi.org/10.1016/j.cpr.2016.07.002>
- Bleichert, J., Wilhelm, F. H., Williams, H., Braams, B. R., Jou, J., & Gross, J. J. (2015). Reappraisal facilitates extinction in healthy and socially anxious individuals. *Journal of Behavior Therapy and Experimental Psychiatry*, 46, 141–150. <https://doi.org/10.1016/j.jbtep.2014.10.001>
- Bottesi, G., Marino, C., Vieno, A., Ghisi, M., & Spada, M. M. (2021). Psychological distress in the context of the COVID-19 pandemic: The joint contribution of intolerance of uncertainty and cyberchondria. *Psychology and Health*, 1–18. <https://doi.org/10.1080/08870446.2021.1952584>
- Britton, J. C., Lissek, S., Grillon, C., Norcross, M. A., & Pine, D. S. (2011). Development of anxiety: The role of threat appraisal and fear learning. *Depression and Anxiety*, 28(1), 5–17. <https://doi.org/10.1002/da.20733>
- Bruine de Bruin, W., Saw, H. W., & Goldman, D. P. (2020). Political polarization in US residents' COVID-19 risk perceptions, policy preferences, and protective behaviors. *Journal of Risk and Uncertainty*, 61(2), 177–194. <https://doi.org/10.1007/s11166-020-09336-3>
- Cannito, L., Di Crosta, A., Palumbo, R., Ceccato, I., Anzani, S., La Malva, P., et al. (2020). Health anxiety and attentional bias toward virus-related stimuli during the COVID-19 pandemic. *Scientific Reports*, 10(1), 1–8. <https://doi.org/10.1038/s41598-020-73599-8>
- Carleton, R. N. (2016). Into the unknown: A review and synthesis of contemporary models involving uncertainty. *Journal of Anxiety Disorders*, 39, 30–43. <https://doi.org/10.1016/j.janxdis.2016.02.007>
- Carr, A. T. (1974). Compulsive neurosis: A review of the literature. *Psychological Bulletin*, 81(5), 311–318. <https://doi.org/10.1037/H0036473>
- Caserotti, M., Girardi, P., Rubaltelli, E., Tasso, A., Lotto, L., & Gavaruzzi, T. (2021). Associations of COVID-19 risk perception with vaccine hesitancy over time for Italian residents. *Social Science & Medicine*, 272, 113688. <https://doi.org/10.1016/j.socscimed.2021.113688>
- Çelik, S., & Köse, G. G. (2021). Mediating effect of intolerance of uncertainty in the relationship between coping styles with stress during pandemic (COVID-19) process and compulsive buying behavior. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 110, 110321. <https://doi.org/10.1016/j.pnpbp.2021.110321>
- Cisler, J. M., & Koster, E. H. W. (2010). Mechanisms of attentional biases towards threat in anxiety disorders: An integrative review. *Clinical Psychology Review*, 30(2), 203–216. <https://doi.org/10.1016/J.CPR.2009.11.003>
- Clark, D. A., & Beck, A. T. (2010). *Cognitive therapy of anxiety disorders: Science and practice*. Guilford Press.
- Coelho, C. M., Suttiwan, P., Arato, N., & Zsido, A. N. (2020). On the nature of fear and anxiety triggered by COVID-19. *Frontiers in Psychology*, 11, 3109. <https://doi.org/10.3389/fpsyg.2020.581314>
- Cohen, R. (2021). *Across the world, covid anxiety and depression take hold*. December 13. New York Times <https://www.nytimes.com/2021/12/13/world/europe/covid-anxiety-depression-omicron.html>.
- Consolidated Appropriations Act. (2021). 133, congress (p. 116). <https://www.congress.gov/bill/116th-congress/house-bill/133/text>.
- Correa, K. A., Liu, H., & Shankman, S. A. (2019). The role of intolerance of uncertainty in current and remitted internalizing and externalizing psychopathology. *Journal of Anxiety Disorders*, 62, 68–76. <https://doi.org/10.1016/J.JANXDIS.2019.01.001>
- Craske, M. G., Hermans, D., & Vervliet, B. (2018). State-of-the-art and future directions for extinction as a translational model for fear and anxiety. *Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences*, 373(1742), 20170025. <https://doi.org/10.1098/rstb.2017.0025>
- Craske, M. G., Kircanski, K., Zelikowsky, M., Mystkowski, J., Chowdhury, N., & Baker, A. (2008). Optimizing inhibitory learning during exposure therapy. *Behaviour Research and Therapy*, 46(1), 5–27. <https://doi.org/10.1016/j.brat.2007.10.003>
- Daniel-Watanabe, L., & Fletcher, P. C. (2021). Are fear and anxiety truly distinct? *Biological Psychiatry Global Open Science*. <https://doi.org/10.1016/j.bpsgos.2021.09.006>
- Davidson, R. J. (1998). Affective style and affective disorders: Perspectives from affective neuroscience. *Cognition & Emotion*, 12(3), 307–330. <https://doi.org/10.1080/026999398379628>
- Denson, T. F., Spanovic, M., & Miller, N. (2009). Cognitive appraisals and emotions predict cortisol and immune responses: A meta-analysis of acute laboratory social stressors and emotion inductions. *Psychological Bulletin*, 135(6), 823–853. <https://doi.org/10.1037/a0016909>
- Doane, L. D., & Adam, E. K. (2010). Loneliness and cortisol: Momentary, day-to-day, and trait associations. *Psychoneuroendocrinology*, 35(3), 430–441. <https://doi.org/10.1016/j.psyneuen.2009.08.005>
- Draheim, A. A., & Anderson, P. L. (2021). Does cognitive behavioral therapy for anxiety disorders improve threat reappraisal?: A meta-analysis. *Journal of Behavioral and Cognitive Therapy*, 31(2), 125–135. <https://doi.org/10.1016/J.JBCT.2020.12.004>
- Dubé, J. P., Smith, M. M., Sherry, S. B., Hewitt, P. L., & Stewart, S. H. (2021). Suicide behaviors during the COVID-19 pandemic: A meta-analysis of 54 studies. *Psychiatry Research*, 301, 113998. <https://doi.org/10.1016/j.psychres.2021.113998>
- Dugas, M. J., & Ladouceur, R. (2000). Treatment of GAD: Targeting intolerance of uncertainty in two types of worry. *Behavior Modification*, 24(5), 635–657. <https://doi.org/10.1177/0145445500245002>
- Dugas, M. J., Letarte, H., Rhéaume, J., Freeston, M. H., & Ladouceur, R. (1995). Worry and problem solving: Evidence of a specific relationship. *Cognitive Therapy and Research*, 19(1), 109–120. <https://doi.org/10.1007/BF02229679>
- Duits, P., Cath, D. C., Lissek, S., Hox, J. J., Hamm, A. O., Engelhard, I. M., et al. (2015). Updated meta-analysis of classical fear conditioning in the anxiety disorders. *Depression and Anxiety*, 32(4), 239–253. <https://doi.org/10.1002/da.22353>
- Ehler, U., Gaab, J., & Heinrichs, M. (2001). Psychoneuroendocrinological contributions to the etiology of depression, posttraumatic stress disorder, and stress-related bodily disorders: The role of the hypothalamus-pituitary-adrenal axis. *Biological Psychology*, 57(1–3), 141–152. [https://doi.org/10.1016/S0301-0511\(01\)00092-8](https://doi.org/10.1016/S0301-0511(01)00092-8)
- Engelhard, I. M., van Uijen, S. L., van Seters, N., & Velu, N. (2015). The effects of safety behavior directed towards a safety cue on perceptions of threat. *Behavior Therapy*, 46(5), 604–610. <https://doi.org/10.1016/j.beth.2014.12.006>
- Etkin, A., & Wager, T. D. (2007). Functional neuroimaging of anxiety: A meta-analysis of emotional processing in PTSD, social anxiety disorder, and specific phobia. *American Journal of Psychiatry*, 164(10), 1476–1488. <https://doi.org/10.1176/appi.ajp.2007.07030504>
- Evans, T. C., Walukevich, K. A., & Britton, J. C. (2016). Vigilance-avoidance and disengagement are differentially associated with fear and avoidant behaviors in social anxiety. *Journal of Affective Disorders*, 199, 124–131. <https://doi.org/10.1016/j.jad.2016.04.003>
- Evans, T. C., Walukevich, K. A., Seager, I., & Britton, J. C. (2018). A psychometric comparison of anxiety-relevant attention measures. *Anxiety, Stress & Coping*, 31(5), 539–554. <https://doi.org/10.1080/10615806.2018.1489536>
- Fergus, T. A. (2013). Cyberchondria and intolerance of uncertainty: Examining when individuals experience health anxiety in response to internet searches for medical information. *Cyberpsychology, Behavior, and Social Networking*, 16(10), 735–739. <https://doi.org/10.1089/cyber.2012.0671>
- Field, A. P. (2006). Is conditioning a useful framework for understanding the development and treatment of phobias? *Clinical Psychology Review*, 26(7), 857–875. <https://doi.org/10.1016/j.cpr.2005.05.010>
- Flores, A., López, F. J., Vervliet, B., & Cobos, P. L. (2020). Prospective intolerance of uncertainty is associated with maladaptive temporal distribution of avoidance responses: An extension of Flores, López, Vervliet, and Cobos (2018). *Journal of Behavior Therapy and Experimental Psychiatry*, 68, 101527. <https://doi.org/10.1016/j.jbtep.2019.101527>
- Foa, E. B., & Kozak, M. J. (1986). Emotional processing of fear: Exposure to corrective information. *Psychological Bulletin*, 99(1), 20–35. <https://doi.org/10.1037/0033-2909.99.1.20>
- Freeston, M., Tiplady, A., Mawn, L., Bottesi, G., & Thwaites, S. (2020). Towards a model of uncertainty distress in the context of Coronavirus (COVID-19). In *Cognitive behaviour therapist* (Vol. 13) Cambridge University Press. <https://doi.org/10.1017/S1754470X2000029X>.

- Fried, E. I., & Robinaugh, D. J. (2020). Systems all the way down: Embracing complexity in mental health research. *BMC Medicine*, 18(1), 1–4. <https://doi.org/10.1186/s12916-020-01668-w>
- Funkhouser, C. J., Correa, K. A., Letkiewicz, A. M., Cozza, E. M., Estabrook, R., & Shankman, S. A. (2021). Evaluating the criterion validity of hierarchical psychopathology dimensions across models: Familial aggregation and associations with Research Domain Criteria (sub)constructs. *Journal of Abnormal Psychology*, 130(6), 575–586. <https://doi.org/10.1037/abn0000687>
- Genes, E. L., & Ruscio, A. M. (2011). A meta-analysis of the relation of intolerance of uncertainty to symptoms of generalized anxiety disorder, major depressive disorder, and obsessive-compulsive disorder. *Clinical Psychology Review*, 31(6), 923–933. <https://doi.org/10.1016/j.cpr.2011.05.001>
- Gloster, A. T., Lamnisos, D., Lubenko, J., Presti, G., Squatrito, V., Constantinou, M., et al. (2020). Impact of COVID-19 pandemic on mental health: An international study. *PLoS One*, 15(12), Article e0244809. <https://doi.org/10.1371/journal.pone.0244809>
- Gorka, S. M., Lieberman, L., Nelson, B. D., Sarapas, C., & Shankman, S. A. (2014). Aversive responding to safety signals in panic disorder: The moderating role of intolerance of uncertainty. *Journal of Anxiety Disorders*, 28(7), 731–736. <https://doi.org/10.1016/j.janxdis.2014.06.004>
- Graupensperger, S., Abdallah, D. A., & Lee, C. M. (2021). Social norms and vaccine uptake: College students' COVID vaccination intentions, attitudes, and estimated peer norms and comparisons with influenza vaccine. *Vaccine*, 39(15), 2060–2067. <https://doi.org/10.1016/j.vaccine.2021.03.018>
- Graupensperger, S., Lee, C. M., & Larimer, M. E. (2021). Young adults underestimate how well peers adhere to COVID-19 preventive behavioral guidelines. *Journal of Primary Prevention*, 42(3), 309–318. <https://doi.org/10.1007/s10935-021-00633-4>
- Greenglass, E., Chiacchia, D., & Fischenbaum, L. (2021). Investigating COVID-19 stress and coping: Substance use and behavioural disengagement. *International Journal of Psychology: Journal International de Psychologie*. <https://doi.org/10.1002/ijop.12820>
- Grillon, C. (2002). Startle reactivity and anxiety disorders: Aversive conditioning, context, and neurobiology. *Biological Psychiatry*, 52(10), 958–975. [https://doi.org/10.1016/S0006-3223\(02\)01665-7](https://doi.org/10.1016/S0006-3223(02)01665-7)
- Grillon, C. (2008). Models and mechanisms of anxiety: Evidence from startle studies. *Psychopharmacology*, 199(3), 421. <https://doi.org/10.1007/s00213-007-1019-1>
- Grupe, D. W., & Nitschke, J. B. (2013). Uncertainty and anticipation in anxiety. *Nature Reviews Neuroscience*, 14(7), 488–501. <https://doi.org/10.1038/nrn3524>
- Harper, C. A., Satchell, L. P., Fido, D., & Litzman, R. D. (2020). Functional fear predicts public health compliance in the COVID-19 pandemic. *International Journal of Mental Health and Addiction*, 1. <https://doi.org/10.1007/S11469-020-00281-5>
- Heeren, A., & McNally, R. J. (2016). An integrative network approach to social anxiety disorder: The complex dynamic interplay among attentional bias for threat, attentional control, and symptoms. *Journal of Anxiety Disorders*, 42. <https://doi.org/10.1016/j.janxdis.2016.06.009>
- van der Heiden, C., Muris, P., & van der Molen, H. T. (2012). Randomized controlled trial on the effectiveness of metacognitive therapy and intolerance-of-uncertainty therapy for generalized anxiety disorder. *Behaviour Research and Therapy*, 50(2), 100–109. <https://doi.org/10.1016/j.brat.2011.12.005>
- Hoffart, A., Johnson, S. U., & Ebrahimi, O. V. (2020). Loneliness and social distancing during the COVID-19 pandemic: Risk factors and associations with psychopathology. *Frontiers in Psychiatry*, 11, 589127. <https://doi.org/10.3389/fpsy.2020.589127>
- Hofmann, S. G., & Hay, A. C. (2018). Rethinking avoidance: Toward a balanced approach to avoidance in treating anxiety disorders. *Journal of Anxiety Disorders*, 55, 14–21. <https://doi.org/10.1016/j.janxdis.2018.03.004>
- Holmes, E. A., O'Connor, R. C., Perry, V. H., Tracey, I., Wessely, S., Arseneault, L., et al. (2020). Multidisciplinary research priorities for the COVID-19 pandemic: A call for action for mental health science. *The Lancet Psychiatry*, 7(6), 547–560. [https://doi.org/10.1016/S2215-0366\(20\)30168-1](https://doi.org/10.1016/S2215-0366(20)30168-1)
- Huang, C., Huang, L., Wang, Y., Li, X., Ren, L., Gu, X., et al. (2021). 6-month consequences of COVID-19 in patients discharged from hospital: A cohort study. *The Lancet*, 397(10270), 220–232. [https://doi.org/10.1016/S0140-6736\(20\)32656-8](https://doi.org/10.1016/S0140-6736(20)32656-8)
- Insel, T. R., Cuthbert, B. N., Garvey, M., Heinssen, R., Pine, D. S., Quinn, K., et al. (2010). Research Domain Criteria (RDoC): Toward a new classification framework for research on mental disorders. *American Journal of Psychiatry*, 167(7), 748–751. <https://doi.org/10.1176/appi.ajp.2010.09091379>
- Jovanovic, T., Kazama, A., Bachevalier, J., & Davis, M. (2012). Impaired safety signal learning may be a biomarker of PTSD. *Neuropharmacology*, 62(2), 695–704. <https://doi.org/10.1016/j.neuropharm.2011.02.023>
- Khakpoor, S., Mohammadi Bytamar, J., & Saed, O. (2019). Reductions in transdiagnostic factors as the potential mechanisms of change in treatment outcomes in the unified protocol: A randomized clinical trial. *Research in Psychotherapy (Milano)*, 22(3), 379. <https://doi.org/10.4081/ripppo.2019.379>
- Khorrami, K. J., Manzler, C. A., Kreutzer, K. A., & Gorka, S. M. (2021). Neural and self-report measures of sensitivity to uncertainty as predictors of COVID-related negative affect. *Psychiatry Research: Neuroimaging*, 319, 111414. <https://doi.org/10.1016/j.psychres.2021.111414>
- Klumpp, H., & Shankman, S. A. (2018). Using event-related potentials and startle to evaluate time course in anxiety and depression. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, 3(1), 10–18. <https://doi.org/10.1016/j.bpsc.2017.09.004>
- Knowles, K. A., & Olatunji, B. O. (2021). Anxiety and safety behavior usage during the COVID-19 pandemic: The prospective role of contamination fear. *Journal of Anxiety Disorders*, 77, 102323. <https://doi.org/10.1016/j.janxdis.2020.102323>
- Koster, E. H. W., Crombez, G., Verschuere, B., & De Houwer, J. (2004). Selective attention to threat in the dot probe paradigm: Differentiating vigilance and difficulty to disengage. *Behaviour Research and Therapy*, 42(10), 1183–1192. <https://doi.org/10.1016/j.brat.2003.08.001>
- Kotov, R., Krueger, R. F., Watson, D., Achenbach, T. M., Althoff, R. R., Bagby, R. M., et al. (2017). The hierarchical taxonomy of psychopathology (HiTOP): A dimensional alternative to traditional nosologies. *Journal of Abnormal Psychology*, 126(4), 454–477. <https://doi.org/10.1037/abn0000258>
- Krypotos, A.-M., Effting, M., Kindt, M., & Beckers, T. (2015). Avoidance learning: A review of theoretical models and recent developments. *Frontiers in Behavioral Neuroscience*, 9, 189. <https://doi.org/10.3389/fnbeh.2015.00189>
- Ladouceur, R., Gosselin, P., & Dugas, M. J. (2000). Experimental manipulation of intolerance of uncertainty: A study of a theoretical model of worry. *Behaviour Research and Therapy*, 38(9), 933–941. [https://doi.org/10.1016/S0005-7967\(99\)00133-3](https://doi.org/10.1016/S0005-7967(99)00133-3)
- Lang, P. J. (2010). Emotion and motivation: Toward consensus definitions and a common research purpose. *Emotion Review*, 2(3), 229–233. <https://doi.org/10.1177/1754073910361984>
- Larsen, E. M., Donaldson, K. R., Liew, M., & Mohanty, A. (2021). Conspiratorial thinking during COVID-19: The roles of paranoia, delusion-proneness, and intolerance of uncertainty. *Frontiers in Psychiatry*, 12, 698147. <https://doi.org/10.3389/fpsy.2021.698147>
- Lazarus, R. S. (1991). Progress on a cognitive-motivational-relational theory of emotion. *American Psychologist*, 46(8), 819–834. <https://doi.org/10.1037/0003-066X.46.8.819>
- Lieberman, L., Funkhouser, C. J., Gorka, S. M., Liu, H., Correa, K. A., Berenz, E. C., et al. (2020). The relation between posttraumatic stress symptom severity and startle potentiation to predictable and unpredictable threat. *The Journal of Nervous and Mental Disease*, 208(5), 397–402. <https://doi.org/10.1097/NMD.0000000000001138>
- Lissek, S., Powers, A. S., McClure, E. B., Phelps, E. A., Woldehawariat, G., Grillon, C., et al. (2005). Classical fear conditioning in the anxiety disorders: A meta-analysis. *Behaviour Research and Therapy*, 43(11), 1391–1424. <https://doi.org/10.1016/j.brat.2004.10.007>
- Lovibond, P. F., Davis, N. R., & O'Flaherty, A. S. (2000). Protection from extinction in human fear conditioning. *Behaviour Research and Therapy*, 38(10), 967–983. [https://doi.org/10.1016/S0005-7967\(99\)00121-7](https://doi.org/10.1016/S0005-7967(99)00121-7)
- Lovibond, P. F., Mitchell, C. J., Minard, E., Brady, A., & Menzies, R. G. (2009). Safety behaviours preserve threat beliefs: Protection from extinction of human fear conditioning by an avoidance response. *Behaviour Research and Therapy*, 47(8), 716–720. <https://doi.org/10.1016/j.brat.2009.04.013>
- MacLeod, C., Mathews, A., & Tata, P. (1986). Attentional bias in emotional disorders. *Journal of Abnormal Psychology*, 95(1), 15–20. <https://doi.org/10.1037/0021-843X.95.1.15>
- Mahoney, A. E. J., & McEvoy, P. M. (2012). Changes in intolerance of uncertainty during cognitive behavior group therapy for social phobia. *Journal of Behavior Therapy and Experimental Psychiatry*, 43(2), 849–854. <https://doi.org/10.1016/j.jbtep.2011.12.004>
- Marek, R., Strobel, C., Bredy, T. W., & Sah, P. (2013). The amygdala and medial prefrontal cortex: Partners in the fear circuit. *The Journal of Physiology*, 591(10), 2381–2391. <https://doi.org/10.1113/jphysiol.2012.248575>
- Martinez-Berman, L., McCutcheon, L., & Huynh, H. P. (2021). Is the worship of celebrities associated with resistance to vaccinations? Relationships between celebrity admiration, anti-vaccination attitudes, and beliefs in conspiracy. *Psychology Health & Medicine*, 26(9), 1063–1072. <https://doi.org/10.1080/13548506.2020.1778754>
- McEvoy, P. M., & Erceg-Hurn, D. M. (2016). The search for universal transdiagnostic and trans-therapy change processes: Evidence for intolerance of uncertainty. *Journal of Anxiety Disorders*, 41, 96–107. <https://doi.org/10.1016/j.janxdis.2016.02.002>
- McEwen, B. S., & Akil, H. (2020). Revisiting the stress concept: Implications for affective disorders. *Journal of Neuroscience*, 40(1), 12–21. <https://doi.org/10.1523/JNEUROSCI.0733-19.2019>
- Meissel, E. E. E., Liu, H., Stevens, E. S., Evans, T. C., Britton, J. C., Letkiewicz, A. M., et al. (2021). The reliability and validity of response-based measures of attention bias. *Cognitive Therapy and Research*, 1–15. <https://doi.org/10.1007/s10608-021-10212-w>
- Mertens, G., Gerritsen, L., Duijndam, S., Saleminck, E., & Engelhard, I. M. (2020). Fear of the coronavirus (COVID-19): Predictors in an online study conducted in March 2020. *Journal of Anxiety Disorders*, 74, 102258. <https://doi.org/10.1016/j.janxdis.2020.102258>
- Mertens, G., & Morris, J. (2021). Intolerance of uncertainty and threat reversal: A conceptual replication of Morris et al. *Behaviour Research and Therapy*, 137, 103799. <https://doi.org/10.1016/j.brat.2020.103799>
- Monninger, M., Pollok, T. M., Aggensteiner, P.-M., Kaiser, A., Reinhard, I., Hermann, A., et al. (2021). Coping under stress: Prefrontal control predicts stress burden during the COVID-19 crisis. *European Neuropsychopharmacology*. <https://doi.org/10.1016/j.euroneuro.2021.11.007>
- Monroe, S. M., & Simons, A. D. (1991). Diathesis-stress theories in the context of life stress research: Implications for the depressive disorders. *Psychological Bulletin*, 110(3), 406–425. <https://doi.org/10.1037/0033-2909.110.3.406>
- Moors, A., Ellsworth, P. C., Scherer, K. R., & Frijda, N. H. (2013). Appraisal theories of emotion: State of the art and future development. *Emotion Review*, 5(2), 119–124. <https://doi.org/10.1177/1754073912468165>
- Morris, J., Bennett, K. P., & Larson, C. L. (2021). I told you it was safe: Associations between intolerance of uncertainty and different parameters of uncertainty during instructed threat of shock. *Journal of Behavior Therapy and Experimental Psychiatry*, 70, 101620. <https://doi.org/10.1016/j.jbtep.2020.101620>

- Morriss, J., & McSorley, E. (2019). Intolerance of uncertainty is associated with reduced attentional inhibition in the absence of direct threat. *Behaviour Research and Therapy*, 118, 1–6. <https://doi.org/10.1016/j.brat.2019.03.011>
- Morriss, J., McSorley, E., & van Reekum, C. M. (2018). I don't know where to look: The impact of intolerance of uncertainty on saccades towards non-predictive emotional face distractors. *Cognition & Emotion*, 32(5), 953–962. <https://doi.org/10.1080/02699931.2017.1370418>
- Morriss, J., & van Reekum, C. M. (2019). I feel safe when i know: Contingency instruction promotes threat extinction in high intolerance of uncertainty individuals. *Behaviour Research and Therapy*, 116, 111–118. <https://doi.org/10.1016/j.brat.2019.03.004>
- Morriss, J., Wake, S., Elizabeth, C., & van Reekum, C. M. (2021). I doubt it is safe: A meta-analysis of self-reported intolerance of uncertainty and threat extinction training. *Biological Psychiatry Global Open Science*, 1(3), 171–179. <https://doi.org/10.1016/j.bpsgos.2021.05.011>
- Morriss, J., Zuj, D. V., & Mertens, G. (2021). The role of intolerance of uncertainty in classical threat conditioning: Recent developments and directions for future research. *International Journal of Psychophysiology: Official Journal of the International Organization of Psychophysiology*, 166, 116–126. <https://doi.org/10.1016/j.ijpsycho.2021.05.011>
- Myers, K. M., & Davis, M. (2007). Mechanisms of fear extinction. *Molecular Psychiatry*, 12(2), 120–150. <https://doi.org/10.1038/sj.mp.4001939>
- Naeem, S. B., & Bhatti, R. (2020). The covid-19 'infodemic': A new front for information professionals. *Health Information and Libraries Journal*, 37(3), 233–239. <https://doi.org/10.1111/hir.12311>
- Nasiry, S., & Ameli, Z. (2021). Effects of online attentional bias modification on coronavirus anxiety. *Practice in Clinical Psychology*, 9(2), 143–152. <https://doi.org/10.32598/jpcp.9.2.739.2>
- Nekliudov, N. A., Blyuss, O., Cheung, K. Y., Petrou, L., Genuneit, J., Sushentsev, N., et al. (2020). Excessive media consumption about COVID-19 is associated with increased state anxiety: Outcomes of a large online survey in Russia. *Journal of Medical Internet Research*, 22(9), Article e20955. <https://doi.org/10.2196/20955>
- Nelson, B. D., Shankman, S. A., & Proudfit, G. H. (2014). Intolerance of uncertainty mediates reduced reward anticipation in major depressive disorder. *Journal of Affective Disorders*, 158, 108–113. <https://doi.org/10.1016/j.jad.2014.02.014>
- Nickerson, R. S. (1998). Confirmation bias: A ubiquitous phenomenon in many guises. *Review of General Psychology*, 2(2), 175–220. <https://doi.org/10.1037/1089-2680.2.2.175>
- Nisa, C. F., Bélanger, J. J., Faller, D. G., Buttrick, N. R., Mierau, J. O., Austin, M. M. K., et al. (2021). Lives versus livelihoods? Perceived economic risk has a stronger association with support for COVID-19 preventive measures than perceived health risk. *Scientific Reports*, 11(1), 1–12. <https://doi.org/10.1038/s41598-021-88314-4>
- Ohman, A., & Mineka, S. (2001). Fears, phobias, and preparedness: Toward an evolved module of fear and fear learning. *Psychological Review*, 108(3), 483–522. <https://doi.org/10.1037/0033-295x.108.3.483>
- Olagoke, A. A., Olagoke, O. O., & Hughes, A. M. (2020). Exposure to coronavirus news on mainstream media: The role of risk perceptions and depression. *British Journal of Health Psychology*, 25(4), 865–874. <https://doi.org/10.1111/bjhp.12427>
- Pepperdine, E., Lomax, C., & Freeston, M. H. (2018). Disentangling intolerance of uncertainty and threat appraisal in everyday situations. *Journal of Anxiety Disorders*, 57, 31–38. <https://doi.org/10.1016/j.janxdis.2018.04.002>
- Pettit, J. W., Bechor, M., Rey, Y., Vasey, M. W., Abend, R., Pine, D. S., et al. (2020). A randomized controlled trial of attention bias modification treatment in youth with treatment-resistant anxiety disorders. *Journal of the American Academy of Child & Adolescent Psychiatry*, 59(1), 157–165. <https://doi.org/10.1016/j.jaac.2019.02.018>
- Presti, G., Mchugh, L., Gloster, A., Karekla, M., & Haye, S. C. (2020). The dynamics of fear at the time of COVID-19: A contextual behavioral science perspective. *Clinical Neuropsychiatry: Journal of Treatment Evaluation*, 17(2), 65–71.
- Rachman, S. (1977). The conditioning theory of fear acquisition: A critical examination. *Behaviour Research and Therapy*, 15(5), 375–387. [https://doi.org/10.1016/0005-7967\(77\)90041-9](https://doi.org/10.1016/0005-7967(77)90041-9)
- Richards, T. N., Nix, J., Mourtgos, S. M., & Adams, I. T. (2021). Comparing 911 and emergency hotline calls for domestic violence in seven cities: What happened when people started staying home due to COVID-19? *Criminology & Public Policy*, 20(3), 573–591. <https://doi.org/10.1111/1745-9133.12564>
- Robinaugh, D., Haslbeck, J., Waldorp, L., Kossakowski, J., Fried, E. I., Millner, A., et al. (2019). Advancing the network theory of mental disorders: A computational model of panic disorder. <https://doi.org/10.31234/osf.io/km37w>
- Robinson, E., Sutin, A. R., Daly, M., & Jones, A. (2021). A systematic review and meta-analysis of longitudinal cohort studies comparing mental health before versus during the COVID-19 pandemic in 2020. *Journal of Affective Disorders*. <https://doi.org/10.1016/j.jad.2021.09.098>
- Roozbeek, J., Schneider, C. R., Dryhurst, S., Kerr, J., Freeman, A. L. J., Recchia, G., et al. (2020). Susceptibility to misinformation about COVID-19 around the world. *Royal Society Open Science*, 7(10). <https://doi.org/10.1098/rsos.201199>
- Rudisill, C. (2013). How do we handle new health risks? Risk perception, optimism, and behaviors regarding the H1N1 virus. *Journal of Risk Research*, 16(8), 959–980. <https://doi.org/10.1080/1080/13669877.2012.761271>
- Sakya, S. M., Scoy, L. J., Van, Garman, J. C., Miller, E. L., Snyder, B., Wasserman, E., et al. (2021). The impact of COVID-19-related changes in media consumption on public knowledge: Results of a cross-sectional survey of Pennsylvania adults. *Current Medical Research and Opinion*, 37(6), 911–915. <https://doi.org/10.1080/03007995.2021.1901679>
- Salkovskis, P. (1996). The cognitive approach to anxiety: Threat beliefs, safety-seeking behavior, and the special case of health anxiety and obsessions. In P. Salkovskis (Ed.), *Frontiers of cognitive therapy* (pp. 48–50). Guildford Press. <https://doi.org/10.23937/2469-5793/1510088>
- San Martín, C., Jacobs, B., & Vervliet, B. (2020). Further characterization of relief dynamics in the conditioning and generalization of avoidance: Effects of distress tolerance and intolerance of uncertainty. *Behaviour Research and Therapy*, 124, 103526. <https://doi.org/10.1016/j.brat.2019.103526>
- Santomauro, D. F., Herrera, A. M. M., Shadid, J., Zheng, P., Ashbaugh, C., Pigott, D. M., et al. (2021). Global prevalence and burden of depressive and anxiety disorders in 204 countries and territories in 2020 due to the COVID-19 pandemic. *The Lancet*, 398(10312), 1700–1712. [https://doi.org/10.1016/S0140-6736\(21\)02143-7](https://doi.org/10.1016/S0140-6736(21)02143-7)
- Satici, B., Saricali, M., Satici, S. A., & Griffiths, M. D. (2020). Intolerance of uncertainty and mental wellbeing: Serial mediation by rumination and fear of COVID-19. *International Journal of Mental Health and Addiction*, 1–12. <https://doi.org/10.1007/s11469-020-00305-0>
- Schlotz, W., Hammerfeld, K., Ehlert, U., & Gaab, J. (2011). Individual differences in the cortisol response to stress in young healthy men: Testing the roles of perceived stress reactivity and threat appraisal using multiphase latent growth curve modeling. *Biological Psychology*, 87(2), 257–264. <https://doi.org/10.1016/j.biopsycho.2011.03.005>
- Schmidt, N. B., Martin, A. D., Allan, N. P., Mathes, B. M., Saulnier, K. G., & March, D. S. (2021). Actual versus perceived infection rates of COVID-19: Impact on distress, behavior and disability. *Journal of Psychiatric Research*, 137, 525–533. <https://doi.org/10.1016/j.jpsychires.2021.03.061>
- Schudy, A., Żurek, K., Wiśniewska, M., Piejka, A., Gawęda, E., & Okruszek, Ł. (2020). Mental well-being during pandemic: The role of cognitive biases and emotion regulation strategies in risk perception and affective response to COVID-19. *Frontiers in Psychiatry*, 11, 589973. <https://doi.org/10.3389/fpsy.2020.589973>
- Shankman, S. A., & Gorka, S. M. (2015). Psychopathology research in the RDoC era: Unanswered questions and the importance of the psychophysiological unit of analysis. *International Journal of Psychophysiology*, 98(2), 330–337. <https://doi.org/10.1016/j.ijpsycho.2015.01.001>
- Shankman, S. A., Gorka, S. M., Nelson, B. D., Fitzgerald, D. A., Phan, K. L., & O'Daly, O. (2014). Anterior insula responds to temporally unpredictable aversiveness: An fMRI study. *NeuroReport*, 25(8), 596–600. <https://doi.org/10.1097/WNR.0000000000000144>
- Sheeran, P., Harris, P. R., & Epton, T. (2014). Does heightening risk appraisals change people's intentions and behavior? A meta-analysis of experimental studies. *Psychological Bulletin*, 140(2), 511–543. <https://doi.org/10.1037/a0033065>
- Shevlin, M., McElroy, E., & Murphy, J. (2015). Loneliness mediates the relationship between childhood trauma and adult psychopathology: Evidence from the adult psychiatric morbidity survey. *Social Psychiatry and Psychiatric Epidemiology*, 50(4), 591–601. <https://doi.org/10.1007/s00127-014-0951-8>
- Skinner, E. A., Edge, K., Altman, J., & Sherwood, H. (2003). Searching for the structure of coping: A review and critique of category systems for classifying ways of coping. *Psychological Bulletin*, 129(2), 216–269. <https://doi.org/10.1037/0033-2909.129.2.216>
- Smith, J. R., Hogg, M. A., Martin, R., & Terry, D. J. (2007). Uncertainty and the influence of group norms in the attitude-behaviour relationship. *British Journal of Social Psychology*, 46(Pt 4), 769–792. <https://doi.org/10.1348/014466606X164439>
- Smits, J. A. J., Julian, K., Rosenfield, D., & Powers, M. B. (2012). Threat reappraisal as a mediator of symptom change in cognitive-behavioral treatment of anxiety disorders: A systematic review. *Journal of Consulting and Clinical Psychology*, 80(4), 624. <https://doi.org/10.1037/A0028957>
- Solís Arce, J. S., Warren, S. S., Meriggi, N. F., Scacco, A., McMurry, N., Voors, M., et al. (2021). COVID-19 vaccine acceptance and hesitancy in low- and middle-income countries. *Nature Medicine*, 27(8), 1385–1394. <https://doi.org/10.1038/s41591-021-01454-y>
- Solomou, I., & Constantinidou, F. (2020). Prevalence and predictors of anxiety and depression symptoms during the COVID-19 pandemic and compliance with precautionary measures: Age and sex matter. *International Journal of Environmental Research and Public Health*, 17(14), 4924. <https://doi.org/10.3390/ijerph17144924>
- Soriano, J. B., Murthy, S., Marshall, J. C., Relan, P., & Diaz, J. V. (2021). A clinical case definition of post-COVID-19 condition by a Delphi consensus. *The Lancet Infectious Diseases*. [https://doi.org/10.1016/S1473-3099\(21\)00703-9](https://doi.org/10.1016/S1473-3099(21)00703-9), 0(0).
- Soroya, S. H., Farooq, A., Mahmood, K., Isoaho, J., & Zara, S. E. (2021). From information seeking to information avoidance: Understanding the health information behavior during a global health crisis. *Information Processing & Management*, 58(2), 102440. <https://doi.org/10.1016/j.ipm.2020.102440>
- Stevens, J. S., Harnett, N. G., Lebois, L. A. M., van Rooij, S. J. H., Ely, T. D., Roekner, A., et al. (2021). Brain-based biotypes of psychiatric vulnerability in the acute aftermath of trauma. *American Journal of Psychiatry*, 178(11), 1037–1049. <https://doi.org/10.1176/appi.ajp.2021.20101526>
- Stokes, P. E. (1995). The potential role of excessive cortisol induced by HPA hyperfunction in the pathogenesis of depression. *European Neuropsychopharmacology: The Journal of the European College of Neuropsychopharmacology*, 5(Suppl), 77–82. [https://doi.org/10.1016/0924-977x\(95\)00039-r](https://doi.org/10.1016/0924-977x(95)00039-r)
- Taha, S., Matheson, K., & Anisman, H. (2014). H1N1 was not all that scary: Uncertainty and stressor appraisals predict anxiety related to a coming viral threat. *Stress and Health*, 30(2), 149–157. <https://doi.org/10.1002/smi.2505>
- Taha, S., Matheson, K., Cronin, T., & Anisman, H. (2014). Intolerance of uncertainty, appraisals, coping, and anxiety: The case of the 2009 H1N1 pandemic. *British Journal of Health Psychology*, 19(3), 592–605. <https://doi.org/10.1111/BJHP.12058>
- Tanovic, E., Gee, D. G., & Joormann, J. (2018). Intolerance of uncertainty: Neural and psychophysiological correlates of the perception of uncertainty as threatening. *Clinical Psychology Review*, 60, 87–99. <https://doi.org/10.1016/j.cpr.2018.01.001>
- Taylor, S. (2021). Understanding and managing pandemic-related panic buying. *Journal of Anxiety Disorders*, 78, 102364. <https://doi.org/10.1016/j.janxdis.2021.102364>

- Taylor, S., Landry, C. A., Paluszek, M. M., & Asmundson, G. J. G. (2020). Reactions to COVID-19: Differential predictors of distress, avoidance, and disregard for social distancing. *Journal of Affective Disorders*, 277, 94–98. <https://doi.org/10.1016/j.jad.2020.08.002>
- Tull, M. T., Barbano, A. C., Scamaldo, K. M., Richmond, J. R., Edmonds, K. A., Rose, J. P., et al. (2020). The prospective influence of COVID-19 affective risk assessments and intolerance of uncertainty on later dimensions of health anxiety. *Journal of Anxiety Disorders*, 75, 102290. <https://doi.org/10.1016/j.janxdis.2020.102290>
- Tversky, A., & Kahneman, D. (1973). Availability: A heuristic for judging frequency and probability. *Cognitive Psychology*, 5(2), 207–232. [https://doi.org/10.1016/0010-0285\(73\)90033-9](https://doi.org/10.1016/0010-0285(73)90033-9)
- van Uijen, S. L., Leer, A., & Engelhard, I. M. (2018). Safety behavior after extinction triggers a return of threat expectancy. *Behavior Therapy*, 49(3), 450–458. <https://doi.org/10.1016/j.beth.2017.08.005>
- Van Damme, S., Crombez, G., Hermans, D., Koster, E. H. W., & Eccleston, C. (2006). The role of extinction and reinstatement in attentional bias to threat: A conditioning approach. *Behaviour Research and Therapy*, 44(11), 1555–1563. <https://doi.org/10.1016/j.brat.2005.11.008>
- Walker, D. L., Toufexis, D. J., & Davis, M. (2003). Role of the bed nucleus of the stria terminalis versus the amygdala in fear, stress, and anxiety. *European Journal of Pharmacology*, 463(1–3), 199–216. [https://doi.org/10.1016/s0014-2999\(03\)01282-2](https://doi.org/10.1016/s0014-2999(03)01282-2)
- Weerakoon, S. M., Jetelina, K. K., & Knell, G. (2021). Longer time spent at home during COVID-19 pandemic is associated with binge drinking among US adults. *The American Journal of Drug and Alcohol Abuse*, 47(1), 98–106. <https://doi.org/10.1080/00952990.2020.1832508>
- Weissman, D. G., Rodman, A. M., Rosen, M. L., Kasperek, S., Mayes, M., Sheridan, M. A., et al. (2021). Contributions of emotion regulation and brain structure and function to adolescent internalizing problems and stress vulnerability during the COVID-19 pandemic: A longitudinal study. *Biological Psychiatry Global Open Science*. <https://doi.org/10.1016/j.bpsgos.2021.06.001>
- Wheaton, M. G., Messner, G. R., & Marks, J. B. (2021). Intolerance of uncertainty as a factor linking obsessive-compulsive symptoms, health anxiety and concerns about the spread of the novel coronavirus (COVID-19) in the United States. *Journal of Obsessive-Compulsive and Related Disorders*, 28, 100605. <https://doi.org/10.1016/j.jocrd.2020.100605>
- White, R. W., & Horvitz, E. (2009). Cyberchondria: Studies of the escalation of medical concerns in Web search. *ACM Transactions on Information Systems*, 10(440), 4. <https://doi.org/10.1145/1629096.1629101>
- Williams, L. M., Goldstein-Piekarski, A. N., Chowdhry, N., Grisanzio, K. A., Haug, N. A., Samara, Z., et al. (2016). Developing a clinical translational neuroscience taxonomy for anxiety and mood disorder: Protocol for the baseline-follow up Research domain criteria Anxiety and Depression (“RAD”) project. *BMC Psychiatry*, 16, 68. <https://doi.org/10.1186/s12888-016-0771-3>
- Woods, A. M., & Bouton, M. E. (2007). Occasional reinforced responses during extinction can slow the rate of reacquisition of an operant response. *Learning and Motivation*, 38(1), 56–74. <https://doi.org/10.1016/j.lmot.2006.07.003>
- Yoon, S., McClean, S. T., Chawla, N., Kim, J. K., Koopman, J., Rosen, C. C., et al. (2021). Working through an “infodemic”: The impact of COVID-19 news consumption on employee uncertainty and work behaviors. *Journal of Applied Psychology*, 106(4), 501–517. <https://doi.org/10.1037/apl0000913>
- Young, K. S., Bookheimer, S. Y., Nusslock, R., Zinbarg, R. E., Damme, K. S. F., Chat, I. K.-Y., et al. (2021). Dysregulation of threat neurocircuitry during fear extinction: The role of anhedonia. *Neuropsychopharmacology: Official Publication of the American College of Neuropsychopharmacology*, 46(9), 1650–1657. <https://doi.org/10.1038/s41386-021-01003-8>
- Zalta, A. K., & Shankman, S. A. (2016). Conducting psychopathology prevention research in the RDoC era. *Clinical Psychology: Science and Practice*, 23(1), 94–104. <https://doi.org/10.1111/cpsp.12144>