



Published in final edited form as:

Int J Methods Psychiatr Res. 2018 March ; 27(1): . doi:10.1002/mpr.1590.

Reliability and Validity of Severity Dimensions of Psychopathology Assessed Using the Structured Clinical Interview for DSM-5 (SCID)

Stewart A. Shankman, Ph.D.^{a,b}, Carter J. Funkhouser, B.A.^a, Daniel N. Klein, Ph.D.^c, Joanne Davila, Ph.D.^c, Debra Lerner, M.S., Ph.D.^d, and Danelle Hee, B.A.^a

^aUniversity of Illinois at Chicago, Department of Psychology

^bUniversity of Illinois at Chicago, Department of Psychiatry

^cStony Brook University, Department of Psychology

^dTufts Medical Center, Program on Work, Health and Productivity, Tufts University School of Medicine and the Sackler Graduate School of Biomedical Sciences, Departments of Medicine and Psychiatry

Abstract

Objective—This study examined whether the Structured Clinical Interview for DSM (SCID), a widely used semi-structured interview designed to assess psychopathology categorically, can be adapted to identify reliable and valid severity dimensions of psychopathology. The present study also examined whether these severity dimensions have better psychometric properties (internal consistency, test-retest reliability, concurrent and predictive validity) than categorical diagnoses.

Methods—Participants (N=234) were recruited from the community and clinics. Retest reliability and prospective predictive validity (symptoms and functioning one-year later) were examined in subsamples of participants. Dimensional severity scales were created from an adapted version of the SCID for both current and lifetime Major Depression, Alcohol, Substance, PTSD, Panic, Agoraphobia, Social Anxiety, Specific Phobia, OCD, and GAD.

Results—The SCID's severity scales demonstrated substantial internal consistency (all Cronbach's alphas >.80), test-retest reliability, concurrent, and predictive validity. Symptom severity scales demonstrated significant incremental validity over and above categorical diagnoses for both current and prospective outcomes.

Conclusions—The psychometric properties of SCID-identified symptom scales were far superior to the psychometrics of categorical diagnoses for both current and lifetime psychopathology. These results highlight the feasibility and utility of the SCID to assess reliable and valid symptom severity dimensions of both current and lifetime psychopathology.

Keywords

SCID; dimensional assessment; reliability; validity

While the modern Diagnostic and Statistical Manuals of Mental Disorders (DSM; post-1980) have provided stakeholders of mental health services (researchers, clinicians, patients, etc.) a common nomenclature to describe psychopathology, the system has received numerous critiques (Thyer, 2015; Wakefield, 2016; Widiger & Clark, 2000). One of the major critiques of the DSM is that it conceptualizes disorders as discrete (i.e., categorical) entities in which hard cut points are used to classify individuals as either “with” or “without” the disorder (Helzer, Kraemer, & Krueger, 2006; Insel et al., 2010). Indeed, numerous taxometric and simulation studies have shown that most psychopathologies are not discrete entities and are better conceptualized as continuous (i.e., dimensional) constructs (Haslam, Holland, & Kuppens, 2012; Ruscio & Marcus, 2007) or perhaps hybrids of dimensional and categorical constructs (Muthen, 2006). Additionally, a meta-analysis reported that compared to when psychopathology is assessed categorically, psychopathology defined dimensionally has 15% greater reliability and 37% greater validity (Markon, Chmielewski, & Miller, 2011).

Despite these psychometric advantages, the best method to assess psychopathology dimensionally is lacking. Most measures designed to assess psychopathology dimensionally are paper-and-pencil questionnaires that either only assess current psychopathology (e.g., a depression scale that assesses symptoms during previous two weeks) or broader traits (e.g., personality disorder measures like the Personality Inventory for DSM5) (Krueger, Derringer, Markon, Watson, & Skodol, 2012). Compared to paper-and-pencil questionnaires, semi-structured interviews such as the Structured Clinical Interview for DSM (SCID; First, Williams, Karg, & Spitzer, 2015) allow for the probing and clarification of responses and are not as impacted by respondents’ reading level (an important concern of many questionnaires (Schinka, 2012)). While there are several interviews designed to assess one or more dimensions of a single area of psychopathology (e.g., Positive and Negative Syndrome Scale for Schizophrenia; Kay, Fiszbein, & Opler, 1987) there is no semi-structured interview designed to assess a broad array of psychopathology both categorically and dimensionally and for both current and past symptoms. Some researchers have used semi-structured interviews like the SCID or Anxiety Disorders Interview Schedule (ADIS-IV; Brown, DiNardo, & Barlow, 1994) to assess psychopathology dimensionally but they have only focused on a single symptom domain (Abela, Skitch, Auerbach, & Adams, 2005) and have not examined past, as well as current, psychopathology (Brown, Chorpita, & Barlow, 1998; Kotov, Gamez, Watson, & Iowa, 2005). Additionally, and most importantly, it is unknown whether these semi-structured interviews yield symptom severity dimensions that are reliable and valid. Having one instrument that can assess an array of psychopathology domains dimensionally would also be useful for studies adopting NIMH’s Research Domain Criteria (RDoC) perspective that encourages a dimensional conceptualization of psychopathology (Sanislow et al., 2010).

The purpose of this study is therefore to (a) adapt a widely used categorical assessment interview of adult psychopathology, the SCID, so that it can also assess the severity of psychopathology dimensionally and (b) compare the reliability and validity of SCID assessed dimensions vs. SCID assessed categories for both current and lifetime psychopathology. We chose to use the SCID for the present study rather than create an

entirely new interview as the SCID is the most widely used semi-structured diagnostic interview for adult psychopathology and its adaptation would not require additional training beyond what SCID users typically receive (SCID-101, 1998).

Method

Participants

Two hundred thirty-four participants were drawn from an ongoing study on neurophysiological vulnerability factors for psychopathology (Gorka et al., 2016; Weinberg, Liu, Hajcak, & Shankman, 2015). The study recruited participants from the community and area mental health clinics (via fliers, Internet postings, etc.) and aimed to obtain a sample of young adults with a wide range of psychiatric diagnoses and symptoms as well as healthy controls. Thus, the inclusion criteria were broad, only requiring age between 18 and 30 years and having at least one full biological sibling within the same age range who was eligible to enroll. Participants were excluded from the larger study if they had a personal or family history of psychosis or mania; were unable to read or write English; had a history of head trauma with loss of consciousness; or were left-handed. These criteria were included to ensure participants were able to provide consent and to mitigate potential confounds to psychophysiological data. Two hundred thirty-four sibling pairs were assessed, but to avoid non-independence of observations, data from only one sibling per family were used for the present analyses. In order to maximize the prevalence of psychopathology, the sibling with the higher sum of lifetime ADIS-IV ratings across the assessed psychopathologies (see below) was selected for the present analyses. Randomly selected subsamples of these 234 individuals were used for the retest reliability ($n=51$) and prospective predictive validity ($n=71$) studies (see below). This investigation was carried out in accordance with the latest version of the Declaration of Helsinki and all participants provided written informed consent after review of the procedures as approved by the university Institutional Review Board. Participants were paid approximately \$20/hour for their participation. Demographic characteristics for these two subsamples and the overall sample are described in Table 1.

Baseline Measures

Structured Clinical Interview for DSM-5—Lifetime and current psychopathology were assessed by interviewers using an adapted version of the research version of the Structured Clinical Interview for DSM-5 (SCID-5-RV; First et al., 2015). Specifically, the following ten disorders were assessed - Major Depressive Disorder (MDD), Alcohol Use Disorder (AUD), Substance Use Disorder (SUD), Post-Traumatic Stress Disorder (PTSD), Panic Disorder (PD), Agoraphobia, Social Anxiety Disorder (SAD), Specific Phobia, Obsessive-Compulsive Disorder (OCD), and Generalized Anxiety Disorder (GAD) (see Supplementary Material for results for the Anorexia, Bulimia, and Binge Eating Disorder modules, which were also administered, but are not reported in the main analyses due to small N). As personal history of psychosis or mania was an exclusion criterion, lifetime psychosis and mania were assessed using the psychotic screening module and manic episode section from the SCID's Mood Disorders module. The SCID for the present study was identical to the SCID-5, with the following exceptions. First, the instrument used a slightly different structure than the SCID-5. The SCID-5 sometimes assesses lifetime diagnostic criteria and

then current criteria for some disorder modules, and sometimes does current and then lifetime (only if the current diagnosis was not met). To aid in the symptom severity assessment of psychopathology, each module of the adapted SCID was organized to always assess lifetime symptoms first and then current symptoms. Second, the separate parts of multicomponent symptoms were coded independently (e.g., the MDD symptom “worthlessness or guilt” [symptom 7] yielded separate ratings for worthlessness *and* guilt). Third, to increase sensitivity to individuals with subthreshold psychopathology and facilitate the calculation of symptom severity scales, we modified some of the skip-out rules in the SCID. Specifically, interviewers ignored all but the first “skip out” for all disorders except MDD and GAD. For example, for PTSD, if a subject received either a 2 or 3 for criterion A (exposure to a trauma), interviewers ignored all subsequent “skip outs” and assessed all lifetime PTSD symptoms even if the subject did not fully meet criteria B, C, D, and/or E (e.g., they only had one symptom from criteria D). However, if a subject received a ‘1’ for criteria A for PTSD (i.e., they never experienced a trauma), the rest of the PTSD symptoms were not assessed as it would not make sense to assess a person’s reactions to a trauma if they never experienced a trauma. Using panic disorder as another example, it would not make sense to assess whether a person worried about additional attacks (Criteria B) if they never had a panic attack (Criteria A). Interviewers also ignored the first “skip out” at the beginning of the MDD and GAD sections because MDD and GAD contain symptoms whose presence are not dependent on the cardinal symptom(s). For example, all of the symptoms of MDD were assessed even if the cardinal symptoms (depressed mood and anhedonia) were not present. Fourth, additional items related to previous course of illness (e.g., duration of longest episode, time since last symptomatic, etc.) were also added to each module (data not included in this report). Fifth, the time frames for current psychopathology were modified for some disorders. Rather than assessing current Agoraphobia, SAD, and Specific Phobia using the past 6 months and assessing current AUD and SUD using the past 12 months (as specified in the SCID-5), we elected to use the past 1 month to assess these current disorders. This modification allowed for the assessment of current symptoms using the same time window as our measures of current functioning (e.g., WHODAS, GAF, and SOFAS), thus facilitating concurrent validity analyses of current psychopathology. Sixth, although the SCID-5 includes questions assessing functional impairment and perceived distress associated with some disorders, several disorders do not include clinically significant impairment or distress as a criterion and do not have questions assessing functional impairment and distress due to symptoms of the specific disorder. Therefore, prompts and rating scales from the ADIS-IV (Brown et al., 1994) were added for each disorder, and for both lifetime and current psychopathology. This provided measures of disorder-specific impairment and distress that were consistent across disorders, thereby facilitating analyses of concurrent validity. These ratings consisted of separate assessments of perceived distress and three forms of impairment (social, occupational, and ‘other’ impairment that resulted from the specific psychopathology) made along a 9-point scale ranging from 0 (*none*) to 8 (*severe*), with (as per ADIS-IV convention) ratings of 2 or higher signifying clinically significant distress or impairment. To provide measures of overall symptom severity and functioning, the Global Assessment of Functioning Scale (“GAF”; American Psychiatric Association., 1994) was modified slightly to only focus on overall symptom severity at the time of the interview and the Social and Occupational Functioning Assessment Scale

(SOFAS; Goldman, Skodol, & Lave, 1992) was used to focus solely on overall functional impairment due to current psychopathology. This is consistent with previous recommendations that the GAF be divided into two scales (Aas, 2010).

Interviewers were trained to criterion by viewing the SCID-101 training videos (SCID-101, 1998), observing two or three SCID interviews with an experienced interviewer, and completing three SCID interviews (observed by the first author or an advanced interviewer) in which diagnoses were in full agreement with those of the observer.

A subset of participants ($n = 51$) completed a second SCID with a different interviewer blind to the first interview within 3 weeks of their first SCID ($M = 8.51$ days, $SD = 4.31$) to assess the test-retest reliability of symptom dimensions and categorical diagnoses.

World Health Organization Disability Assessment Schedule 2.0—To assess more global disability and impairment, the 36-item World Health Organization Disability Assessment Schedule 2.0 (WHODAS; Ustün et al., 2010) interview was administered immediately after the SCID. The 36-item WHODAS is the gold standard assessment of disability published by the World Health Organization and contains six domains of current functioning (Cognition, Mobility, Self-care, Getting along, Life activities, and Participation), which are calculated by summing the values for the items in each domain. These domain level scores have previously shown excellent internal consistency and test-retest reliability, as well as concurrent, construct, and discriminant validity (Ustün et al., 2010). In the present study, Cronbach's alphas for the WHODAS domains ranged from .81–.92.

One-year Follow-up Assessment

A subset of participants ($n=71$) completed the Longitudinal Interval Follow-up Evaluation interview (LIFE, the gold standard assessment of longitudinal symptom assessment) and 12-item WHODAS interview one year after their initial SCID interview (data was only available for a subset as data collection is ongoing). The 12-item WHODAS interview retrospectively assessed impairment since baseline. It accounts for 81% of the variance of the 36-item version (Ustün et al., 2010), but only produces an overall disability score and not domain level scores. WHODAS impairment ratings were obtained for each month between the baseline assessment and the follow-up.

The LIFE (Keller et al., 1987) interview was administered face-to-face ($n=65$) or over the phone ($n=6$) at one year follow-up to retrospectively assess symptoms of psychopathology since baseline. For each week since baseline, the LIFE yields a measure of symptom severity called a psychiatric status rating (PSR) for each disorder assessed at baseline, ranging from 1 (*absent*) to 6 (*severe*), with ratings of 5 or 6 signifying that the full diagnostic criteria for the disorder are met. The LIFE has exhibited good to excellent inter-rater reliability and excellent test-retest reliability over a one year period (Keller et al., 1987; Warshaw, Keller, & Stout, 1994).

Data Analysis

We measured lifetime and current psychopathology using both categorical diagnoses and symptom severity scales. After all interviews were completed, symptom severity scales were

electronically computed by summing each symptom within a disorder, with each symptom rated on a 1–3 scale (except for the percentage time symptom from OCD). The severity scales included the following symptoms:¹ MDD - nine core symptoms; AUD and SUD - 11 core symptoms; PD - symptom A, B1, and B2; PTSD - symptom A, the 20 symptoms in criterion B through E; Agoraphobia - the five potentially feared situations under symptom A and symptom B, C, D, and E; SAD and specific phobia (scored separately) - symptom A, B, C, D, and E; OCD - symptom A1, A2, B1, B2, and the percentage of time spent engaging in OCD thoughts or behaviors (all z-scored); GAD – symptom A, B, and the six symptoms under symptom C; AN – symptom A, B, and C; BN – symptom A1, A2, B, C, and D; BED – symptom A1 and A2, the five symptoms under criterion B, and symptoms C and D.

Internal consistencies of each symptom severity scale were determined using Cronbach's alphas. One-way random single-measure intraclass correlation coefficients (ICCs) were used to assess the test-retest reliability of severity scales, and Cohen's kappas were used to assess test-retest reliability of categorical diagnoses. For these reliability analyses, we used Shrout's (1998) standards for determining strength of reliabilities (although these standards are not without controversy (Kraemer, Kupfer, Clarke, Narrow, & Regier, 2012); specifically, 0.00 – 0.10 (virtually none), 0.11 – 0.40 (slight), 0.41 – 0.60 (fair), 0.61– 0.80 (moderate), 0.81–1.0 (substantial). For these analyses, there were nine sibling pairs in the test-retest reliability subsample (but not the baseline [$n=234$] or predictive validity [$N=71$] sample). The inclusion of sibling pairs in the test-retest reliability subsample infringes on the statistical assumption that observations are independent. However, results were comparable in a smaller sample using only one sibling from each pair.

Concurrent validity for lifetime and current psychopathology was assessed using correlations between each diagnosis or symptom severity scale (point-biserial and Pearson correlations, respectively) and the corresponding ADIS ratings, with the idea that assessments of psychopathology should be associated with functional impairment. Concurrent validity was also assessed with the domain level scores of the 36-item WHODAS and the GAF current symptom severity and SOFAS current impairment ratings.² The key analyses regarding concurrent validity tested incremental validity of symptom dimensions over and above categorical diagnoses. To test this, we computed multiple regressions with both symptom severity scales and categorical diagnoses as simultaneous predictors of the various validators (ADIS ratings, etc).

The predictive validity of each diagnosis and severity scale was assessed via (1) the maximum LIFE severity rating for the corresponding disorder over the one year follow-up, and (2) the overall impairment index from the 12-item WHODAS across the follow-up (specifically, the average of the maximum rating of each of the 12 items across the follow-up). These measures were used to measure longitudinal risk with the assumption that SCID

¹We also explored the reliability and validity of dimensions if the multiple components of symptoms were included separately (e.g., separate symptoms of worthlessness and guilt for symptom 7 of MDD). The pattern of results with these dimensions was nearly identical to when these components were collapsed. To increase the utility and applicability of the results to labs that do not 'unpack' these symptoms, only results with the 'packed' symptoms are reported below.

²The Mobility and Self-care domains of the 36-item WHODAS were excluded due to a lack of variability in scores (i.e., most participants reported no impairment in these domains).

assessments should be associated with longitudinal risk for psychopathology. Although the relationships between SCID assessments and the maximum LIFE severity rating are likely influenced by the persistence of each disorder, the specific disorders' persistence were equated in the key analyses that examined whether dimensional and categorical assessments (entered simultaneously in a multiple regression) contributed incremental validity over and above each other.

Results

Table 2 presents the internal consistency and test-retest reliability for categorical and dimensional severity assessments of psychopathology. Using Shrout's (Shrout, 1998) conventions for judging reliability coefficients, the internal consistencies for the 20 dimensional assessments of lifetime and current psychopathologies were all in the substantial range (above .80) with the sole exception of current alcohol use disorder (Cronbach's alpha = .78 [moderate range]). For the test-retest reliabilities, all 10 of the dimensional assessments for lifetime psychopathology were in the substantial range, while Kappas for categorical diagnoses demonstrated poorer reliability (2 were substantial, 4 were moderate, 2 were fair, and 1 was slight). A similar pattern emerged for test-retest reliability of current psychopathology with nine of the ten dimensional severity assessments demonstrating substantial reliability (the tenth was moderate). Test-retest reliability was poorer for current categorical diagnoses (none were substantial, 4 were moderate, 2 were fair, and 1 was slight).³

Table 3 shows that both dimensional severity scales and categorical diagnoses of lifetime psychopathology were associated with significant concurrent validity for all validators. Table 3 also shows the results for multiple regression models in which both dimensional severity and categorical assessments of psychopathology were predictors. The symptom severity scales remained significant for 29 of the 30 regression models, while the categorical assessment was only significant for 8 of the 30 models. Additionally, symptom severity scales had larger independent effects than categorical diagnoses for 27 of the 30 models. There was a similar pattern for the concurrent validity of the current psychopathology assessment (see Table 4), albeit slightly weaker compared to lifetime psychopathology given the lower levels of current psychopathology. Specifically, severity scales remained significant for 66 of the 90 (73%) multiple regression models where both dimension severity and categorical assessments were entered simultaneously, compared to only 15 of the categorical effects (15 of 90 [16.7%]). Dimensional assessments had larger effects than categorical diagnoses for 84 of the 90 models (93%).

Table 5 presents the predictive validity of categorical and dimensional severity assessments. Categorical and dimensional assessments for lifetime and current psychopathology prospectively predicted higher symptoms of the corresponding disorder over the course of the 1-year follow-up. Table 5 also shows the results for multiple regression models in which both dimensional and categorical assessments were included. Dimensional severity assessments demonstrated stronger incremental predictive validity for 17 of 20 lifetime

³Analyses for several categorical diagnoses could not be calculated due to small N (see Tables for details).

psychopathology models and 16 of 18 current psychopathology models. Fewer symptom severity scales and categorical diagnoses prospectively predicted WHODAS functioning. However, when both dimensional severity and categorical assessments were included in the same model, only dimensional (and never categorical) assessments demonstrated significant incremental predictive validity for functioning.

Discussion

The present study found that symptom severity dimensions of psychopathology identified by the SCID demonstrate substantial internal consistency, test-retest-reliability, concurrent, and predictive validity, and far superior psychometric properties than categorical diagnoses for most disorders. While studies of personality disorders and research utilizing paper-and-pencil questionnaires have previously demonstrated similarly superior psychometrics for dimensional assessments of psychopathology (Markon et al., 2011), this is the first study to our knowledge that has examined the ability of a semi-structured interview to identify reliable severity dimensions of MDD, GAD, OCD, PTSD, Panic Disorder, Agoraphobia, Social Anxiety, Specific Phobia, alcohol use disorders, and substance use disorders for both current and past psychopathology.

These results suggest that if research labs are seeking to use a semi-structured interview to assess psychopathology dimensionally (or with a categorical/dimensional hybrid model, Muthen, 2006), they can simply adapt the SCID (one of the most widely adopted tools to assess psychopathology) to do so, without additional training beyond the standard SCID. The SCID might be a particularly useful measure for labs conducting RDoC inspired research that define psychopathology dimensionally, but may also need to classify participants' categorical diagnostic status (e.g., for exclusion criteria; Cuthbert & Kozak, 2013).

There are, of course, other measures to assess current psychopathology dimensionally (e.g., Kotov et al., 2005; Watson et al., 2008), but most of these are self-report (which have limitations as discussed above) or do not assess a broad range of psychopathology. Another advantage of the SCID over these instruments is that it not only assesses current (i.e., last several weeks or month) but also lifetime psychopathology and the present results suggest that severity dimensions for both can be assessed reliably and validly. It is interesting that symptom severity scales of lifetime psychopathology were reliable and valid given questions about the validity of categorically defined lifetime psychopathology assessments (Olino et al., 2012). These prior studies did not assess lifetime psychopathology dimensionally, however, and this may have contributed to their poorer validity.

We do not assume that the DSM-based severity scales used in this study provide the most parsimonious classification system. We took a 'top-down' approach to defining severity dimensions, using the structure of the SCID (which is organized by DSM categories), while most studies examining the structure of psychopathology take a 'bottom up' approach, examining how symptoms of psychopathology co-vary into parsimonious dimensions and/or types (Eaton, Krueger, Docherty, & Sponheim, 2014). It will be important for future studies to develop semi-structured interviews that measure 'higher-order' dimensions (such as the

frequently observed dimensions of internalizing and externalizing psychopathology; Farmer, Seeley, Kosty, Olino, & Lewinsohn, 2013; Krueger & Markon, 2006).

Strengths of the study included a well-characterized sample of individuals with broad range of symptomatology, a longitudinal follow-up, re-test assessments with a different interviewer, and numerous measures of functioning to validate the measures of psychopathology. There were also several notable limitations. First, individuals with a personal or family history of bipolar disorders or psychosis were excluded, thus limiting the study's ability to examine those psychopathologies. Second, impairment was reported by the respondent and was not corroborated by independent report, although interviewers attempted to elicit examples from respondents in order to validate these ratings during the interview. Third, several disorders that were assessed had very few cases for the categorical analyses (particularly for the analyses of current psychopathology). Fourth, the present study only examined the dimension of symptom severity and not other dimensions that are important to psychopathology assessment (e.g., duration of symptoms; Shankman & Klein, 2002). Fifth, the sample was not systematically recruited for the purposes of this study and had a limited age range (18–30), which limits the generalizability of these findings to other clinical or community samples or other age groups.

In sum, the present study suggests that even though the SCID was designed to assess psychopathology categorically, it can be adapted to yield severity dimensions of the common forms of psychopathology (depression, anxiety, substance and alcohol use) that are reliable and valid. Most importantly, these dimensions consistently showed incremental validity over and above SCID-defined categories of psychopathology. As the field considers moving towards frameworks that define psychopathology dimensionally (Helzer et al., 2006), additional studies such as the present one are valuable in assessing the psychometrics (and ultimately feasibility and utility) of dimensional assessment approaches.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding Sources

This work was supported by a grant from the National Institute of Mental Health (R01MH098093 [PI: Shankman]) and the National Center for Advancing Translational Sciences, National Institutes of Health (UL1TR002003).

References

- Aas IH. Global Assessment of Functioning (GAF): properties and frontier of current knowledge. *Annals of General Psychiatry*. 2010; 9:20.doi: 10.1186/1744-859X-9-20 [PubMed: 20459646]
- Abela JR, Skitch SA, Auerbach RP, Adams P. The impact of parental borderline personality disorder on vulnerability to depression in children of affectively ill parents. *Journal of Personality Disorders*. 2005; 19(1):68–83. DOI: 10.1521/pedi.19.1.68.62177 [PubMed: 15899721]
- American Psychiatric Association. *Diagnostic and statistical manual of mental disorders : DSM-IV*. 4. Washington, DC: American Psychiatric Association; 1994.

- Brown TA, Chorpita BF, Barlow DH. Structural relationships among dimensions of the DSM-IV anxiety and mood disorders and dimensions of negative affect, positive affect, and autonomic arousal. *Journal of Abnormal Psychology*. 1998; 107(2):179–192. [PubMed: 9604548]
- Brown, TA., DiNardo, P., Barlow, DH. *Anxiety Disorder Interview Schedule (ADIS-IV & ADIS-IV-L)*. Graywind Publications Inc; 1994.
- Cuthbert BN, Kozak MJ. Constructing constructs for psychopathology: the NIMH research domain criteria. *Journal of Abnormal Psychology*. 2013; 122(3):928–937. [PubMed: 24016027]
- Eaton NR, Krueger RF, Docherty AR, Sponheim SR. Toward a model-based approach to the clinical assessment of personality psychopathology. *Journal of Personality Assessment*. 2014; 96:283–292. [PubMed: 24007309]
- Farmer RF, Seeley JR, Kosty DB, Olino TM, Lewinsohn PM. Hierarchical organization of axis I psychiatric disorder comorbidity through age 30. *Comprehensive Psychiatry*. 2013; 54(5):523–532. DOI: 10.1016/j.comppsy.2012.12.007 [PubMed: 23332721]
- First, MB., Williams, JBW., Karg, RS., Spitzer, RL. *Structured Clinical Interview for DSM-5: Research Version*. Arlington, VA: American Psychiatric Association; 2015.
- Goldman HH, Skodol AE, Lave TR. Revising axis V for DSM-IV: a review of measures of social functioning. *American Journal of Psychiatry*. 1992; 149(9):1148–1156. DOI: 10.1176/ajp.149.9.1148 [PubMed: 1386964]
- Gorka SM, Hee D, Lieberman L, Mittal VA, Phan KL, Shankman SA. Reactivity to uncertain threat as a familial vulnerability factor for alcohol use disorder. *Psychological Medicine*. 2016; :1–10. DOI: 10.1017/s0033291716002415
- Haslam N, Holland E, Kuppens P. Categories versus dimensions in personality and psychopathology: a quantitative review of taxometric research. *Psychological Medicine*. 2012; 42(5):903–920. DOI: 10.1017/s0033291711001966 [PubMed: 21939592]
- Helzer JE, Kraemer HC, Krueger RF. The feasibility and need for dimensional psychiatric diagnoses. *Psychological Medicine*. 2006; 36(12):1671–1680. [PubMed: 16907995]
- Insel T, Cuthbert B, Garvey M, Heinssen R, Pine DS, Quinn K, ... Wang P. Research domain criteria (RDoC): toward a new classification framework for research on mental disorders. *American Journal of Psychiatry*. 2010; 167(7):748–751. [PubMed: 20595427]
- Kay SR, Fiszbein A, Opler LA. The positive and negative syndrome scale (PANSS) for schizophrenia. *Schizophrenia Bulletin*. 1987; 13(2):261–276.
- Keller MB, Lavori PW, Friedman B, Nielsen E, Endicott J, McDonald-Scott P, Andreasen NC. The Longitudinal Interval Follow-up Evaluation. A comprehensive method for assessing outcome in prospective longitudinal studies. *Archives of General Psychiatry*. 1987; 44(6):540–548. [PubMed: 3579500]
- Kotov, R., Gamez, W., Watson, D. Unpublished manuscript. University of Iowa; 2005. Development and validation of the Interview for Mood and Anxiety Symptoms (IMAS).
- Kraemer HC, Kupfer DJ, Clarke DE, Narrow WE, Regier DA. DSM-5: How reliable is reliable enough? *American Journal of Psychiatry*. 2012; 169:13–15. [PubMed: 22223009]
- Krueger RF, Derringer J, Markon KE, Watson D, Skodol AE. Initial construction of a maladaptive personality trait model and inventory for DSM-5. *Psychological Medicine*. 2012; 42(9):1879–1890. [PubMed: 22153017]
- Krueger RF, Markon KE. Reinterpreting comorbidity: a model-based approach to understanding and classifying psychopathology. *Annual Review of Clinical Psychology*. 2006; 2:111–133.
- Markon KE, Chmielewski M, Miller CJ. The reliability and validity of discrete and continuous measures of psychopathology: a quantitative review. *Psychological Bulletin*. 2011; 137(5):856–879. DOI: 10.1037/a0023678 [PubMed: 21574681]
- Muthen B. Should substance use disorders be considered as categorical or dimensional? *Addiction*. 2006; 101:6–16.
- Olino T, Shankman S, Klein D, Seeley J, Pettit J, Farmer R, Lewinsohn P. Lifetime rates of psychopathology in single versus multiple diagnostic assessments: Comparison in a community sample of probands and siblings. *Journal of Psychiatric Research*. 2012; 46(9):1217–1222. DOI: 10.1016/j.jpsy.2012.05.017 [PubMed: 22739001]

- Regier DA, Narrow WE, Clarke DE, Kraemer HC, Kuramoto SJ, Kuhl EA, Kupfer DJ. DSM-5 field trials in the United States and Canada, Part II: test-retest reliability of selected categorical diagnoses. *American Journal of Psychiatry*. 2013; 170(1):59–70. DOI: 10.1176/appi.ajp.2012.12070999 [PubMed: 23111466]
- Ruscio J, Marcus DK. Detecting small taxa using simulated comparison data: A reanalysis of Beach, Amir, and Bau's (2005) data. *Psychological Assessment*. 2007; 19(2):241–246. DOI: 10.1037/1040-3590.19.2.241 [PubMed: 17563206]
- Sanislow CA, Pine DS, Quinn KJ, Kozak MJ, Garvey MA, Heihsen RK, ... Cuthbert BN. Developing constructs for psychopathology research: research domain criteria. *Journal of Abnormal Psychology*. 2010; 119(4):631–639. [PubMed: 20939653]
- Schinka JA. Further issues in determining the readability of self-report items: comment on McHugh and Behar (2009). *Journal of Consulting and Clinical Psychology*. 2012; 80(5):952–955. DOI: 10.1037/a0029928 [PubMed: 22924333]
- SCID-101 (Writer). SCID-101 for DSM-IV: Training video for the Structured Clinical Interview for DSM-(SCID) [Motion Picture]. N. Y. S. P. I. Biometrics Research Department (Producer); 1998.
- Shankman S, Klein D. Dimensional diagnosis of depression: Adding the dimension of course to severity, and comparison to the DSM. *Comprehensive Psychiatry*. 2002; 43(6):420–426. DOI: 10.1053/comp.2002.35902 [PubMed: 12439827]
- Shrout PE. Measurement reliability and agreement in psychiatry. *Stat Methods Med Res*. 1998; 7(3): 301–317. [PubMed: 9803527]
- Thyer, BA. The DSM-5 definition of mental disorder: Critique and alternatives. In: Probst, B., editor. *Critical thinking in clinical assessment and diagnosis*. Cham, Switzerland: Springer; 2015. p. 45-68.
- Ustün TB, Chatterji S, Kostanjsek N, Rehm J, Kennedy C, Epping-Jordan J. ... Project, W. N. J. Developing the World Health Organization Disability Assessment Schedule 2.0. *Bulletin of the World Health Organization*. 2010; 88(11):815–823. [PubMed: 21076562]
- Warshaw MG, Keller MB, Stout RL. Reliability and validity of the longitudinal interval follow-up evaluation for assessing outcome of anxiety disorders. *Journal of Psychiatric Research*. 1994; 28(6):531–545. [PubMed: 7699612]
- Watson D, O'Hara MW, Chmielewski M, McDade-Montez EA, Koffel E, Naragon K, Stuart S. Further validation of the IDAS: evidence of convergent, discriminant, criterion, and incremental validity. *Psychological Assessment*. 2008; 20(3):248–259. [PubMed: 18778161]
- Wakefield JC. Diagnostic issues and controversies in DSM-5: Return of the false positives problem. *Annual Review of Clinical Psychology*. 2016; 12:105–132.
- Weinberg A, Liu H, Hajcak G, Shankman SA. Blunted neural response to rewards as a vulnerability factor for depression: Results from a family study. *Journal of Abnormal Psychology*. 2015; 124(4): 878–889. DOI: 10.1037/abn0000081 [PubMed: 26214708]
- Widiger TA, Clark LA. Toward DSM—V and the classification of psychopathology. *Psychological Bulletin*. 2000; 126(6):946–963. [PubMed: 11107884]

Table 1

Sample Characteristics

	Whole Sample (N=234)	Test-Retest Sample (N=51)	Predictive Validity Sample (N=71)
Age (<i>M, SD</i>)	22.46 (3.19)	22.27 (3.20)	22.23 (3.36)
Gender			
Male	76 (32.5%)	21 (41.2%)	24 (33.8%)
Female	158 (67.5%)	30 (58.8%)	47 (66.2%)
Ethnicity			
Caucasian	96 (41.0%)	21 (41.2%)	29 (40.8%)
Hispanic	53 (22.6%)	6 (11.8%)	19 (26.8%)
African American	34 (14.5%)	12 (23.5%)	9 (12.7%)
Asian	25 (10.7%)	5 (9.8%)	8 (11.3%)
Middle Eastern	10 (4.3%)	1 (2.0%)	2 (2.8%)
Mixed Race	14 (6.0%)	5 (9.8%)	3 (4.2%)
Other	2 (.9%)	1 (2.0%)	1 (1.4%)
Mean GAF (<i>M, SD</i>)			
Symptoms	68.06 (14.00)	67.44 (10.88)	68.38 (16.04)
Impairment	70.80 (13.36)	70.54 (10.77)	70.85 (14.69)
Education			
Did not graduate high school	10 (4.2%)	1 (2.0%)	2 (2.8%)
Graduated high school (or equivalent)	10 (4.3%)	6 (11.8%)	2 (2.8%)
Some College	120 (51.3%)	24 (47.1%)	44 (62.0%)
Graduated 2 year college	12 (5.1%)	5 (9.8%)	0 (.0%)
Graduated 4 year college	43 (18.4%)	11 (21.6%)	15 (21.1%)
Some graduate or professional school	28 (12.0%)	4 (7.8%)	5 (7.0%)
Completed graduate or professional school	10 (4.3%)	0 (.0%)	3 (4.2%)
Employment Status			
Full-time	59 (25.2%)	10 (19.6%)	19 (26.8%)
Part-time	55 (23.5%)	11 (21.6%)	23 (32.4%)
Student	106 (45.3%)	24 (47.1%)	26 (36.6%)
Unemployed	14 (6.0%)	6 (11.8%)	3 (4.2%)
Lifetime Psychiatric Hospitalization	17 (7.3%)	6 (11.8%)	8 (11.3%)
Lifetime Psychiatric Treatment	99 (42.3%)	20 (39.2%)	32 (45.1%)

Table 2

Internal consistency (n=234) and test-retest reliability of current and lifetime psychopathology (N=51)

	Cronbach's Alpha (n=234)	Test-Retest Reliability (ICCs [95% CI] for Sx severity scales; kappas [95% CI] for categorical Dx - n=51)	# of Participants Diagnosed in Time 1	# of Participants Diagnosed in Time 2
Lifetime MDD				
Categorical Dx	-	k = .69 [.49, .89]	26	28
Sx Severity	.91	.91 ** [.84, .95]		
Current MDD				
Categorical Dx	-	k = .74 [.46, 1.00]	8	5
Sx Severity	.91	.85 ** [.76, .91]		
Lifetime AUD				
Categorical Dx	-	k = .84 [.68, .99]	21	19
Sx Severity	.88	.79 ** [.76, .91]		
Current AUD				
Categorical Dx	-	k = .54 [.08, 1.00]	3	4
Sx Severity	.78	.38 ** [.13, .59]		
Lifetime SUD				
Categorical Dx	-	k = .87 [.73, 1.00]	19	18
Sx Severity	.91	.92 ** [.87, .96]		
Current SUD				
Categorical Dx	-	k = .64 [.26, 1.00]	5	4
Sx Severity	.89	.76 ** [.61, .85]		
Lifetime PTSD				
Categorical Dx	-	k = .65 [.20, 1.00]	2	4
Sx Severity	.94	.85 ** [.75, .91]		
Current PTSD				
Categorical Dx	-	can't be computed because one set had no one diagnosed	1	0
Sx Severity	.89	.73 ** [.57, .84]		
Lifetime Panic				
Categorical Dx	-	k = .46 [.12, .79]	9	6
Sx Severity	.86	.74 ** [.58, .84]		
Current Panic				
Categorical Dx	-	can't be computed because one set had no one diagnosed	1	0
Sx Severity	.85	.12 [-.16, .38]		
Lifetime Agoraphobia				
Categorical Dx	-	can't be computed because one set had no one diagnosed	2	0
Sx Severity	.91	.63 ** [.44, .77]		
Current Agoraphobia				

	Cronbach's Alpha (n=234)	Test-Retest Reliability (ICCs [95% CI] for Sx severity scales; kappas [95% CI] for categorical Dx - n=51)	# of Participants Diagnosed in Time 1	# of Participants Diagnosed in Time 2
Categorical Dx	-	can't be computed because one set had no one diagnosed	1	0
Sx Severity	.92	.31* [.04, .53]		
Lifetime Social Anxiety				
Categorical Dx	-	k = .18 [-.15, .43]	13	14
Sx Severity	.97	.61** [.40, .75]		
Current Social Anxiety				
Categorical Dx	-	k = .29 [-.10, .68]	6	5
Sx Severity	.98	.58** [.36, .73]		
Lifetime Specific Phobia				
Categorical Dx	-	k = .71 [.44, .97]	10	6
Sx Severity	.97	.72** [.55, .83]		
Current Specific Phobia				
Categorical Dx	-	k = .70 [.38, 1.00]	7	4
Sx Severity	.97	.77** [.63, .86]		
Lifetime OCD				
Categorical Dx	-	k = .49 [.11, .88]	5	6
Sx Severity	.92	.61** [.40, .76]		
Current OCD				
Categorical Dx	-	k = .64 [.26, 1.00]	4	5
Sx Severity	.88	.57** [.35, .73]		
Lifetime GAD				
Categorical Dx	-	k = .77 [.51, 1.00]	7	8
Sx Severity	.94	.79** [.67, .88]		
Current GAD				
Categorical Dx	-	k = .56 [.18, .94]	4	6
Sx Severity	.95	.76** [.61, .85]		

Note. Sx = symptoms; Dx = Diagnosis MDD = Major Depressive Disorder, AUD = Alcohol Use Disorder; SUD = Substance Use Disorder, PTSD = Post-traumatic stress disorder, OCD = Obsessive-Compulsive Disorder, GAD = Generalized anxiety disorder.

* $p < .05$.

** $p < .01$.

Table 3Concurrent validity for lifetime psychopathology (N=234)

	Lifetime ADIS social	Lifetime ADIS occupational	Lifetime ADIS distress
Lifetime MDD			
Categorical Dx (prev = 46.2%)	-.02 (.66 ^{**})	-.09 (.58 ^{**})	-.12 ⁺ (.64 ^{**})
Sx Severity	.83 ^{**} (.81 ^{**})	.82 ^{**} (.74 ^{**})	.92 ^{**} (.82 ^{**})
Lifetime AUD			
Categorical Dx (prev = 32.1%)	-.14 ⁺ (.52 ^{**})	.18 [*] (.69 ^{**})	-.12 (.48 ^{**})
Sx Severity	.80 ^{**} (.69 ^{**})	.62 ^{**} (.77 ^{**})	.73 ^{**} (.63 ^{**})
Lifetime SUD			
Categorical Dx (prev = 26.1%)	.04 (.64 ^{**})	.00 (.65 ^{**})	.04 (.54 ^{**})
Sx Severity	.66 ^{**} (.71 ^{**})	.73 ^{**} (.73 ^{**})	.57 ^{**} (.60 ^{**})
Lifetime PTSD			
Categorical Dx (prev = 12.0%)	.00 (.63 ^{**})	.04 (.57 ^{**})	.03 (.63 ^{**})
Sx Severity	.83 ^{**} (.83 ^{**})	.70 ^{**} (.72 ^{**})	.78 ^{**} (.81 ^{**})
Lifetime Panic			
Categorical Dx (prev = 11.1%)	-.03 (.69 ^{**})	.03 (.66 ^{**})	-.01 (.73 ^{**})
Sx Severity	.84 ^{**} (.82 ^{**})	.73 ^{**} (.76 ^{**})	.87 ^{**} (.86 ^{**})
Lifetime Agoraphobia			
Categorical Dx (prev = 2.1%)	-.01 (.52 ^{**})	-.03 (.46 ^{**})	-.05 (.52 ^{**})
Sx Severity	.85 ^{**} (.84 ^{**})	.79 ^{**} (.77 ^{**})	.92 ^{**} (.89 ^{**})
Lifetime Social Anxiety			
Categorical Dx (prev = 21.4%)	.18 ^{**} (.56 ^{**})	.11 (.52 ^{**})	.08 ⁺ (.61 ^{**})
Sx Severity	.56 ^{**} (.68 ^{**})	.62 ^{**} (.69 ^{**})	.77 ^{**} (.83 ^{**})
Lifetime Specific Phobia			
Categorical Dx (prev = 23.5%)	.24 ^{**} (.47 ^{**})	.38 ^{**} (.44 ^{**})	.36 ^{**} (.70 ^{**})
Sx Severity	.32 ^{**} (.49 ^{**})	.08 (.36 ^{**})	.47 ^{**} (.73 ^{**})
Lifetime OCD			
Categorical Dx (prev = 9.8%)	.41 ^{**} (.66 ^{**})	.08 (.48 ^{**})	.48 ^{**} (.75 ^{**})
Sx Severity	.32 ^{**} (.64 ^{**})	.50 ^{**} (.57 ^{**})	.33 ^{**} (.72 ^{**})
Lifetime GAD			
Categorical Dx (prev = 14.5%)	-.11 [*] (.47 ^{**})	-.07 (.50 ^{**})	-.14 ^{**} (.48 ^{**})
Sx Severity	.89 ^{**} (.82 ^{**})	.87 ^{**} (.82 ^{**})	.96 ^{**} (.87 ^{**})

Note. In each cell, zero-order correlations are presented inside the bracket and standardized betas from multiple regressions are presented outside of bracket. Betas are adjusted for other assessment (i.e., categorical adjusted for severity, and severity adjusted for categorical). Cat = categorical, prev = prevalence, MDD = Major Depressive Disorder, AUD = Alcohol Use Disorder, SUD = Substance Use Disorder, PTSD = Post-traumatic stress disorder, OCD = Obsessive-Compulsive Disorder, ADIS = Anxiety Disorders Interview Schedule.

⁺ p<.10,

^{*} p<.05,

**
p<.01,

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 4

Concurrent validity for current psychopathology (N=234)

	Current ADIS social	Current ADIS occupational	Current ADIS distress	GAF Symptoms	SOEAS Impairment	WHODAS cognition	WHODAS getting along	WHODAS life activities	WHODAS participation
Current MDD									
Cat. Dx (prev = 8.1%)	-.41** (.50**)	-.38** (.49**)	-.31** (.57**)	.19 ⁺ (-.31**)	.21* (-.31**)	-.08 (.35**)	-.20 ⁺ (.23**)	-.23* (.27**)	-.19 ⁺ (.33**)
Sx Sev.	1.13** (.80**)	1.08** (.77**)	1.09** (.84**)	-.62** (-.46**)	-.64** (-.47**)	.53** (.46**)	.54** (.38**)	.63** (.44**)	.65** (.50**)
Current AUD									
Cat. Dx (prev = 6.0%)	-.03 (.38**)	.44** (.55**)	-.26** (.47**)	-.07 (-.12 ⁺)	.01 (-.10)	.07 (.07)	-.06 (.07)	-.12 (.07)	-.06 (.03)
Sx Sev.	.48** (.45**)	.13 (.51**)	.85** (.63**)	-.06 (-.12 ⁺)	-.13 (-.12 ⁺)	-.01 (.06)	.15 (.10)	.22 ⁺ (.12 ⁺)	.10 (.05)
Current SUD									
Cat. Dx (prev = 6.8%)	-.01 (.53**)	.02 (.63**)	-.14 (.51**)	.08 (-.16**)	.13 (-.13**)	.02 (.13 ⁺)	-.24 ⁺ (.09)	-.28* (.06)	-.26* (.00)
Sx Sev.	.64** (.63**)	.70** (.72**)	.75** (.63**)	-.38** (-.29**)	-.30* (-.20**)	.13 (.14*)	.38** (.18**)	.40** (.16*)	.30* (.08)
Current PTSD									
Cat. Dx (prev = 2.1%)	.10 ⁺ (.52**)	.06 (.40**)	-.22** (.31**)	.04 (-.20**)	.10 (-.16*)	-.06 (.05)	-.06 (.05)	-.04 (.14*)	.01 (.15*)
Sx Sev.	.66** (.73**)	.55** (.58**)	.84** (.70**)	-.35** (-.30**)	-.40** (-.34**)	.17* (.13*)	.18* (.14*)	.28** (.26**)	.22** (.23**)
Current Panic									
Cat. Dx (prev = 3.8%)	.16** (.74**)	.41** (.82**)	.16* (.69**)	.19 ⁺ (-.24**)	.14 (-.22**)	-.01 (.12 ⁺)	.06 (.21**)	.12 (.20**)	.09 (.23*)
Sx Sev.	.72** (.84**)	.51** (.84**)	.66** (.79**)	-.52** (-.37**)	-.45** (-.32**)	.17 (.16*)	.18 (.23*)	.10 (.19**)	.17 (.24**)
Current Agoraphobia									
Cat. Dx (prev = 1.3%)	-.07 (.43**)	-.24** (.29**)	-.16** (.42**)	.14 ⁺ (-.10)	.10 (-.12 ⁺)	-.09 (.07)	-.10 (.08)	.00 (.06)	.06 (.12 ⁺)
Sx Sev.	.84** (.80**)	.89** (.74**)	.97** (.88**)	-.41** (-.32**)	-.36** (-.31**)	.26** (.21**)	.30** (.24**)	.10 (.10)	.10 (.14*)
Current Social Anxiety									
Cat. Dx (prev = 12.0%)	.15* (.56*)	.15* (.57**)	.18** (.66**)	-.05 (-.25**)	-.14 ⁺ (-.32**)	.07 (.19**)	.00 (.25**)	-.03 (.09)	-.06 (.11 ⁺)
Sx Sev.	.62** (.72**)	.62** (.72**)	.73** (.84**)	-.30** (-.34**)	-.27** (-.36**)	.18* (.23**)	.37** (.28**)	.17 ⁺ (.16*)	.26** (.22**)
Current Spec. Phobia									
Cat. Dx (prev = 16.2%)	.29** (.49**)	.31** (.47**)	.39** (.69**)	-.19* (-.32**)	-.14 (-.26*)	.16 ⁺ (.18**)	.13 (.17*)	.19* (.18**)	.13 (.14*)
Sx Sev.	.29** (.49**)	.23** (.44**)	.43** (.70**)	-.18* (-.31**)	-.17 ⁺ (-.27**)	.03 (.14*)	.05 (.14*)	-.01 (.12 ⁺)	.01 (.10)
Current OCD									

	Current ADIS social	Current ADIS occupational	Current ADIS distress	GAF Symptoms	SOFAS Impairment	WHODAS cognition	WHODAS getting along	WHODAS life activities	WHODAS participation
Cat. Dx (prev = 6.8%)	.62** (.40**)	.46** (.07)	.64** (.22**)	-.25** (-.04)	-.20** (-.01)	.04 (-.04)	.11 (.04)	.13 ⁺ (.04)	.16* (.12 ⁺)
Sx Sev.	.51** (.05)	.53** (.30**)	.68** (.36**)	-.29** (-.14*)	-.25** (-.15*)	.08 (.07)	.11 (.04)	.14* (.05)	.10 (-.03)
Current GAD									
Cat. Dx (prev = 6.4%)	-.08 ⁺ (.47**)	.02 (.55**)	-.09* (.50**)	.01 (-.28**)	.05 (-.25**)	.01 (.27**)	.13 ⁺ (.32**)	-.04 (-.18**)	.04 (.29**)
Sx Sev.	.89** (.84**)	.87** (.88**)	.96** (.91**)	-.49** (-.48**)	-.49** (-.46**)	.42** (.42**)	.32** (.40**)	.35** (.33**)	.41** (.43**)

Note. In each cell, zero-order correlations are presented inside the bracket and betas adjusted for other assessment are presented outside the bracket (i.e., categorical adjusted for severity, and severity adjusted for categorical). Cat = categorical; Sev = severity; prev = prevalence, MDD = Major Depressive Disorder, AUD = Alcohol Use Disorder; SUD = Substance Use Disorder, PTSD = Post-traumatic stress disorder, OCD = Obsessive-Compulsive Disorder, WHODAS = World Health Organization Disability Assessment Scale.

⁺ p<.10,

* p<.05,

** p<.01,

Table 5

Current and Lifetime Psychopathology Predicting Symptoms and Functioning at 1-year follow-up (predictive validity)

	Max PSR over 1 year	WHODAS (mean _{max}) at 1 year FU
Lifetime MDD		
Categorical Dx (prev. = 45.1%)	.12 (.48 **)	-.19 (.28 *)
Dimensional Sx	.44* (.54 **)	.57** (.42 **)
Current MDD		
Categorical Dx (prev. = 8.5%)	-.11 (.40 **)	-.01 (.31 **)
Sx Severity	.62** (.54 **)	.41* (.40 **)
Lifetime AUD		
Categorical Dx (prev. = 35.2%)	-.09 (.32 **)	-.20 (-.11)
Sx Severity	.40* (.42 **)	.11 (-.06)
Current AUD		
Categorical Dx (prev. = 9.9%)	.27 (.54 **)	.08 (-.07)
Sx Severity	.31 (.54 **)	-.18 (-.10)
Lifetime SUD		
Categorical Dx (prev. = 23.9%)	-.05 (.57 **)	-.08 (.03)
Sx Severity	.69** (.65 **)	.12 (.05)
Current SUD		
Categorical Dx (prev. = 11.3%)	.22 (.64 **)	.24 (.01)
Sx Severity	.49** (.68 **)	-.27 (-.06)
Lifetime PTSD		
Categorical Dx (prev. = 12.7%)	.25 ⁺ (.62 **)	.34 ⁺ (.25 *)
Sx Severity	.47** (.67 **)	-.12 (.14)
Current PTSD		
Categorical Dx (prev. = 2.8%)	-.08 (.43 **)	-.18 (-.02)
Sx Severity	.81** (.76 **)	.26 (.14)
Lifetime Panic		
Categorical Dx (prev. = 12.7%)	-.16 (.38 **)	-.26 (.25 *)
Sx Severity	.63** (.49 **)	.61** (.38 **)
Current Panic		
Categorical Dx (prev. = 4.2%)	-.14 (.45 **)	-.01 (.04)
Sx Severity	.73** (.62 **)	.06 (.05)
Lifetime Agoraphobia		
Categorical Dx (prev. = 1.4%)	-.37** (-.01)	-.21 (-.05)
Sx Severity	.58** (.35 **)	.26 ⁺ (.13)
Current Agoraphobia		

	Max PSR over 1 year	WHODAS (mean _{max}) at 1 year FU
Categorical Dx (prev. = 0.0%)	N/A	N/A
Sx Severity	N/A (.54 ^{**})	N/A (-.03)
Lifetime Social Anxiety		
Categorical Dx (prev. = 18.3%)	.40 ^{**} (.65 ^{**})	.08 (.23 ⁺)
Sx Severity	.37 ^{**} (.65 ^{**})	.23 (.28 [*])
Current Social Anxiety		
Categorical Dx (prev. = 9.9%)	.11 (.63 ^{**})	.07 (.28 [*])
Sx Severity	.77 ^{**} (.85 ^{**})	.31 [*] (.36 ^{**})
Lifetime Specific Phobia		
Categorical Dx (prev. = 21.1%)	.22 (.52 ^{**})	-.03 (.10)
Sx Severity	.41 ^{**} (.57 ^{**})	.17 (.15)
Current Specific Phobia		
Categorical Dx (prev. = 15.5%)	.29 ^{**} (.66 ^{**})	.10 (.09)
Sx Severity	.54 ^{**} (.74 ^{**})	-.02 (.05)
Lifetime OCD		
Categorical Dx (prev. = 12.7%)	-.18 (.48 ^{**})	.12 (.36 ^{**})
Sx Severity	.83 ^{**} (.68 ^{**})	.31 (.40 ^{**})
Current OCD		
Categorical Dx (prev. = 11.3%)	-.06 (.51 ^{**})	.24 (.29 [*])
Sx Severity	.72 ^{**} (.67 ^{**})	.06 (.26 [*])
Lifetime GAD		
Categorical Dx (prev. = 14.1%)	-.02 (.46 ^{**})	.09 (.27 [*])
Sx Severity	.73 ^{**} (.72 ^{**})	.28 ⁺ (.34 ^{**})
Current GAD		
Categorical Dx (prev. = 9.9%)	.02 (.48 ^{**})	.13 (.31 ^{**})
Sx Severity	.76 ^{**} (.77 ^{**})	.31 [*] (.39 ^{**})

Note. In each cell, zero-order correlations are presented inside of bracket and betas adjusted for the other assessment are presented outside the bracket (i.e., categorical adjusted for severity, and severity adjusted for categorical). Dx = Diagnosis, Sx = symptoms, prev = prevalence, MDD = Major Depressive Disorder, AUD = Alcohol Use Disorder; SUD = Substance Use Disorder, PTSD = Post-traumatic stress disorder, OCD = Obsessive-Compulsive Disorder, PSR = Psychiatric Status Rating (from the LIFE interview) for the psychopathology in the respective row, WHODAS = World Health Organization Disability Assessment Scale.

⁺ p<.10,

^{*} p<.05,

^{**} p<.01,